

United States  
Department of  
Agriculture

Forest Service

Technology &  
Development  
Program

2400-Timber  
February 1996  
9624-2808-MTDC



# Materials and Supplies for Management of Wildlife Damage to Trees



# Materials and Supplies for Management of Wildlife Damage to Trees



Dale L. Nolte, *Wildlife Biologist*

Ivy J. Otto, *Wildlife Technician*

**USDA Animal and Plant Health Inspection Service  
Animal Damage Control  
Denver Wildlife Research Center  
Olympia, Washington**

**USDA Forest Service  
Technology & Development Program  
Missoula, Montana**

**4E42E33—Animal Repellents**

**February 1996**

---

The Forest Service, United States Department of Agriculture, has developed this information for the guidance of its employees, its contractors, and its cooperating Federal and State agencies, and is not responsible for the interpretation or use of this information by anyone except its own employees. The use of trade, firm, or corporation names in this publication is for the information and convenience of the reader, and does not constitute an endorsement by the Department of any product or service to the exclusion of others that may be suitable.

The United States Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means of communication of program information (braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-2791. To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call (202) 720-7327 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.

This publication reports research involving pesticides. It does not contain recommendation for their use, nor does it imply that the uses discussed here have been registered. All uses of pesticides must be registered by appropriate State and/or Federal agencies before they can be recommended. CAUTION: Pesticides can be injurious to humans, domestic animals, desirable plants, and fish or other wildlife—if they are not handled or applied properly. Use all pesticides selectively and carefully. Follow recommended practices for the disposal of surplus pesticide and pesticide containers.

Check all Federal and State laws and regulations before implementing a damage prevention program. The applicator assumes responsibility to ensure that all legal and safety concerns are met prior to using products. Carefully read and follow the restrictions and safety precautions provided on applicable labels. Federal and State registrations certify that it is legal to use the product according to the conditions and restriction stipulated on the approved label. Registration does not guarantee the availability or the efficacy of a product.

# Contents

Foreword	1
Acknowledgments	2
Introduction	3
Developing a Damage Reduction Program	4
Problem Identification	5
Mountain Beaver ( <i>Aplodontia rufa</i> )	5
Beaver ( <i>Castor canadensis</i> )	5
Elk ( <i>Cervus</i> spp.)	6
Porcupine ( <i>Erethizon dorsatum</i> )	6
Snowshoe Hare ( <i>Lepus americanus</i> )	7
Vole ( <i>Microtus</i> spp.)	7
Deer ( <i>Odocoileus</i> spp.)	8
Pocket Gopher ( <i>Thomomys</i> spp.)	8
Black Bear ( <i>Ursus americanus</i> )	9
Physical Deterrents	10
Individual Barriers	10
Tubes	10
Conical Protectors	12
Bud Caps	12
Obstructions	12
Squirrel Guards	12
Exclosures	12
Fencing	12
Netting	15

# Contents

<b>Traps</b>	<b>16</b>
<b>Live Traps</b>	<b>16</b>
Box Traps	16
Suitcase or Clamshell Traps	17
Leg-Hold Traps	18
Foot Snares	19
<b>Lethal Traps</b>	<b>20</b>
Body-Grip Traps	20
Snap Traps	21
Pocket Gopher Traps	22
Neck Snares	22
<b>Toxicants</b>	<b>24</b>
<b>Ingested Toxicants</b>	<b>24</b>
Acute Toxicants	24
Chronic Toxicants	25
<b>Fumigants</b>	<b>26</b>
<b>Applicator Equipment</b>	<b>26</b>
Broadcast Applicators	26
Burrow-Builders	27
Bait Stations	27
Probes and Bait Dispensers	28
<b>Repellents (Chemosensory)</b>	<b>29</b>
Area Repellents	30
Contact Repellents	30

# Contents

<b>Frightening Devices</b>	<b>32</b>
<b>Visual Devices</b>	<b>32</b>
Reflectors	32
Artificial Light	32
Effigies	33
Balloons, Kites, and Flags	33
<b>Noisemakers</b>	<b>34</b>
Distress Calls, Sirens, and Horns	34
Ultrasonic Devices	34
Automatic Exploders	35
Pyrotechnic Devices	36
<b>Alternative Forage</b>	<b>37</b>
Supplemental Bear Feeding Program	37
<b>Sources</b>	<b>39</b>
<b>Bibliography</b>	<b>46</b>



## Foreword

---

**T**he growth and survival of seedlings planted on National Forests has improved dramatically over the past 20 years, primarily because nurseries produce better stock, and because more care is used in handling and planting seedlings.

One remaining problem is that of animal damage to planted stock. Although livestock, rodents, and other animals take their toll, deer and elk are the primary animals browsing seedlings. By nipping off buds and shoots, deer and elk can restrict seedling growth and cause seedling mortality. Fencing plantations can greatly reduce the damage, but the cost is high. Tubes are

sometimes used to protect seedlings, but they are costly and require annual maintenance to be effective. Chemical sprays and powders have been used with varying degrees of success.

Better techniques and products to protect newly planted seedlings from animal damage need to be found. This would reduce reforestation costs, improve seedling survival and growth, and help land managers meet stocking requirements on schedule. The first step in solving this problem is to determine what techniques and products are now being used to protect seedlings or are available to be used.

This catalog is the first step in working toward a solution of the problem of animal damage to newly planted tree seedlings. It documents the various methods and products now available. The Missoula Technology and Development Center contracted with Dr. Dale Nolte of the Animal and Plant Health Inspection Service Animal Damage Control's Denver Wildlife Research Center to prepare this catalog. The USDA Forest Service's Washington Office Timber Management Reforestation Program funded the work.

**Ben Lowman**  
*Program Manager*  
*Missoula Technology and Development Center*

## Acknowledgments

---

The authors would like to acknowledge their appreciation for the comments and suggestions on drafts of this publication offered by Patricia Archuleta, Michael Fall, James

Farley, Lynwood Fiedler, Constance Harrington, Larry Johnson, Ted Otto, Bob Phillips, John Theade, Tim Veenendaal, and Georg Ziegltrum. Pieter Straub gathered information on

sources of materials, contacted companies, and collected product information. Most of the illustrations were drawn by Phil Lundberg from photos and descriptions provided by the authors.

---



## Introduction

---

**F**oraging wildlife can inflict severe damage to forest resources. Any tree is susceptible to attack. However, seedlings are the most vulnerable. Reforestation after a timber harvest or forest fire is often impeded by foraging herbivores. Seedlings are clipped or gnawed by rodents and the foliage is stripped by browsing ungulates. The resultant damage can be lethal or can impair growth rates and induce deformities. Animals also girdle established trees to feed on sapwood. Complete girdling is lethal, while partial girdling slows growth rates and provides avenues for subsequent insect or disease infestations.

A variety of available methods to reduce wildlife damage to forest resources are presented. Each approach and operation are briefly described and most are accompanied by an illustration. Advantages and disadvantages are also given, along with a list of sources to purchase materials.

The **Problem Identification** section very briefly describes identifying characteristics of selected mammals commonly associated with forest damage. The **Physical Deterrent** section describes barriers and fencing options. The section on **Traps** provides information on materials for the removal of animals. The **Toxicants** section presents means to reduce or maintain low population

densities of animals. Chemicals to reduce foraging are provided in the **Repellents** section. A section on **Frightening Devices** lists available visual aids and noisemakers to scare animals away from protected sites. The **Alternative Forage** section describes an approach to alleviate damage by providing wildlife with an alternative to eating trees. A section on **Sources** lists the addresses, phone numbers, and facsimile numbers of the manufacturers and distributors of the products described. Finally, the **Bibliography** provides sources used to prepare this manuscript and more detailed information on tools and methods to protect forest resources from damage by wildlife.





# Developing a Damage Reduction Program

---

**T**he most appropriate approach to reduce animal damage needs to reflect the overall objectives of the producer, as well as the conditions of the specific problem. All techniques are not feasible or appropriate for all situations. The process described below provides a reasonable approach to develop a program to reduce damage.

First, assess the severity, as well as the potential for additional damage to occur if no action is taken. Concurrently, identify the correct culprit, or the target species of your anticipated program.

Second, evaluate the feasibility of all possible approaches to alleviate the problem. No action may be the appropriate action if you decide the problem is relatively minor. Check on the legal ramifications for any action. Then evaluate all legal approaches relative to the biological, environmental, economical and sociocultural aspects

of the problem. Ascertain that your selected action will not be potentially hazardous to endangered or threatened species. An effective approach also requires that you are familiar with behavioral traits and biology of the target species. Evaluate how the environmental conditions of the site may effect your approach and the consequences of your action to the environment. Consider the likelihood that your selected method will achieve an acceptable degree of success, and whether the situation warrants the anticipated expense. Public attitudes toward potential methods also need to be considered when choosing an approach.

Third, develop a strategy to implement your efforts to reduce damage. Your strategy may incorporate several methods at once, or utilize one method to stop the damage and another to limit future problems. Inquire among experts

within the field if you need additional information or are unsure of specific requirements. Acquire training or expertise in handling equipment or chemicals, if necessary. Identify and obtain any required equipment, personnel, resources, and safety equipment for your program.

Fourth, implement your program. Unanticipated problems or concerns may force you to alter strategies. In that case, repeat the decision process incorporating the new information.

Fifth, monitor consequences of your program. Determine whether your desired goals are being achieved and whether there are any unexpected negative consequences. Continue to evaluate the program until the resource is no longer vulnerable, or conditions warrant terminating the program.

## Problem Identification

A critical step in any program to successfully reduce forest resource depredation is to correctly identify the individual animal or species inflicting the damage. Animals presented below are generally recognized as species capable of inflicting severe damage to reforestation efforts or timber crops. These species are common in the forest and almost everyone can recognize them on sight. Unfortunately, the culprit is rarely seen. Therefore, the offending animal may need to be identified solely on the basis of the resultant damage. A brief description of the type of damage inflicted by each species is presented.

### Mountain Beaver (*Aplodontia rufa*)

Mountain beavers clip seedlings and tree branches up to an inch in diameter (Figure 1). Their diagonal cut is typical of rodents, but multiple bites may create a serrated edge. Larger trees suffer basal barking and undermining of roots. Girdling by mountain beavers can be readily distinguished from bear girdling because the damage is lower on the bole and mountain beavers leave horizontal tooth marks and irregular claw marks. Further, bark strips will not be found lying at the base of the tree. Burrow systems of mountain beavers are usually present. Fresh digging, or fresh vegetation and debris near burrow entrances, indicate an active system.

### Beaver (*Castor canadensis*)

Beavers may be destructive to waterways and forest resources. Flooding of roads and timber stands from blocked waterways is generally more detrimental than tree cutting. Tree cutting and barking, however, has

increased substantially during the last few years in the Pacific Northwest. Conical-shaped stumps and large wood chips at the base of stumps are good

indicators of beaver damage (Figure 2). Peeled sticks with uniform horizontal toothmarks are also generally found in the vicinity of beaver activity.




Figure 1—A mountain beaver and its characteristic signs.




Figure 2—A beaver and its characteristic signs.

## Snowshoe Hare (*Lepus americanus*)

Snowshoe hare damage is difficult to distinguish from mountain beaver damage. An oblique, 45° angle cut is generally present on clipped seedlings (Figure 5). Snowshoe hares tend to prefer feeding on seedlings less than 1/4 inch in diameter. The most conspicuous signs of snowshoe hare activity are their tracks and fecal pellets left throughout the damaged site.




Figure 5—A snowshoe hare and its characteristic signs.

## Vole (*Microtus* spp.)

Vole damage, like damage caused by most other forest rodents, occurs predominately on young seedlings. Pointed stems on clipped seedlings, and small whorled or circular marks on girdled seedlings are characteristic on seedlings clipped and girdled by voles (Figure 6). Root damage by voles is recognizable because voles generally strip the root and leave pointed tips. Where voles are active, there are distinct trails and intermittent open burrow holes.




Figure 6—A vole and its characteristic signs.

## Deer (*Odocoileus* spp.)

Deer damage inflicted on seedlings is similar to elk damage. Woody stems are often splintered and the bark is stripped from twigs (Figure 7). New buds are generally clipped back to the previous year's growth in the spring. Deer do not pull seedlings as frequently as elk and their damage rarely occurs above 6 feet.




Figure 7—A deer and its characteristic signs.

## Pocket Gopher (*Thomomys* spp.)

Pocket gophers commonly prune seedling roots, and girdle or clip seedling stems. Small seedlings, less than 1/2 inch in diameter, are the most vulnerable. The stems are clipped at or near ground level (Figure 8). Pocket gophers may pull the harvested seedling into their burrows. Pocket gophers also prune the roots and girdle the stems of larger trees. Extensive aboveground girdling of larger trees occurs under deep snow conditions. Aboveground girdling is fairly easy to detect. Damage to roots, however, may go unnoticed until seedlings tip over or become discolored. Nonlethal damage causes poor overall growth, shortened needles, reduced internodes, premature needle drop, and needle discoloration.




Figure 8—A pocket gopher and its characteristic signs.

## Elk (*Cervus* spp.)

Big game species, such as elk (Figure 3) and deer, inflict the most widespread form of damage to forest resources. Elk may pull newly planted seedlings or seedlings without well-established root systems out of the ground. Browsing elk often splinter woody stems. During the spring, the stems may be stripped of bark below the break. Elk also trample seedlings.




Figure 3—An elk and its characteristic signs.

## Porcupine (*Erethizon dorsatum*)

Porcupines usually forage on young to pole-size trees. However, they may damage trees of all ages. Porcupines strip bark and clip branches of young trees (Figure 4), or forage among the tops of older trees. Branches are generally clipped at a 45° angle, with horizontal and blunt tooth marks (1/8 inch) along stripped branches. Prime indicators of porcupine activity are bark chips, clipped needles, quills, and fecal matter at the base of damaged trees. Porcupines may also girdle the base of trees. During the winter, porcupines leave noticeable trails in the snow as they travel between trees.




Figure 4—A porcupine and its characteristic signs.

## Black Bear (*Ursus americanus*)

Bear damage occurs in the spring as bears become active and foraging opportunities are limited. Timber stands between 15 and 30 years of age are the most commonly damaged. Bears tend to select the more productive trees. Therefore, silvicultural practices that generate growth (e.g., thinning, fertilizing) may attract foraging bears. Bears generally feed on the lower bole of a tree (Figure 9), but occasionally an entire tree is stripped. They strip the bark with their claws and then feed on the sapwood by scraping it from the heartwood with their teeth. Trees damaged by bears have stripped bark lying at their bases. Vertical tooth and claw marks are generally visible.




Figure 9—A bear and its characteristic signs.

# Physical Deterrents

Physical deterrents impede animal access to specific trees or entry into reforestation plantations or nurseries. Deterrents vary from minor efforts to extensive construction projects. The most feasible approach depends on several economic, physical, and biological factors. The value of the resources to be protected, as well as the availability of resources to install a deterrent, also need to be considered.

The species and behavioral characteristics of the animal(s) inflicting the damage will dictate the physical traits of a deterrent. For example, a short compact barrier may protect a seedling from pocket gopher damage but provide no protection against deer browsing or antler rubbing. Conversely, an extensive fence may deter deer, yet fail to reduce damage by pocket gophers. Consideration of seedling stem form may require that a barrier be removed or that it degrades within a specified time frame. Consideration should also be given to any negative effects a physical deterrent may have on nontarget species or on nondamaging behaviors of the target species. Aesthetics may be a concern in some areas.

The most common methods of excluding animals are individual barriers, encircling an area with a fence, or covering resources with netting. Individual barriers are placed around each seedling or used to cover specific plant parts (e.g., terminal bud). A variety of tubes, bud caps, squirrel guards and obstructions are commercially available or easily constructed. Fences are constructed to prevent entry of foraging animals into specified areas. Drift fences are also used to alter animal movements away from a site or to encourage animals to move toward another control measure (i.e., direct porcupines along a fence into a box trap). Netting can be strung between posts to create a temporary fence or used as a cover for seedbeds to alleviate animal predation in nurseries.

Generally, physical deterrents are not hazardous to wildlife or humans. Prior to fence construction, the general area should be evaluated to ensure that physical barriers will not disrupt normal migratory patterns or exclude wildlife from limited resources, such as water or salt. Loose wires or openings in which wildlife could be ensnared need to be eliminated. Likewise, bird netting can be installed inside of tubes to prevent birds from entering and possibly being trapped in tall tubes. These considerations will not only decrease chances for detrimental effects to wildlife, but also greatly increase the longevity and efficacy of the structure.

## Individual Barriers

### Tubes

**Function:** Entire seedlings or portions of their root systems or stems are surrounded by tubes to reduce herbivory by foraging wildlife (Figure 10).

**Description:** A variety of tubing styles are available. Vexar, a plastic mesh made from polypropylene or a polypropylene and polyethylene mixture, was one of the first tubes commercially available.

Currently, tubes are produced from a variety of materials, including cardboard, metal, paper, and plastics. The rate of degradation of a tube will depend largely on the type of material from which it is constructed. Both tube color and placement (e.g., shade versus sunlight) will affect the photodegradation rate of plastic protectors. Dark colors, however, tend to degrade more slowly than light colors. The goal is to select materials that are strong and last long enough to provide the desired protection but degrade before they interfere with tree growth.

The microclimate of seedlings may be altered by tubes. Seedlings within tubes will experience less wind than seedlings planted in the open. The restricted space and protective cover may also cause an increase in temperature, relative humidity, and soil moisture within the tubes. The amount and balance of solar radiation may also be altered.

The altered microclimate may have positive, neutral, or negative effects on the seedling's growth and health depending on the tube type, seedling size and condition, and environmental conditions. Less wind will reduce evaporation losses but will also favor height growth over stem girth, creating an unfavorable elongated seedling. Under cool climatic conditions, temperature increase in the tube will enhance growth. However, increased temperatures will suppress growth if the temperature becomes too hot, and may reduce hardening off during the fall. Increased relative humidity decreases evaporation losses, but also increases the likelihood of foliar diseases. Likewise, increased soil moisture favors growth, but too much soil moisture increases the seedlings susceptibility to diseases. Radiation, dependent on the tube, may be altered to a more favorable light balance to enhance photosynthesis or the tubes may create a less favorable light balance or permit insufficient light for photosynthesis.

Tube walls may be solid or a mesh with diamond, twill, or square patterns. The twill and square patterns are flexible and easy to compress for easy transport. The rigidity of the diamond mesh and solid tubes enables them to withstand more abuse by animals. Solid tubes (Figure 11), or tubes with finer mesh sizes, reduce the likelihood that terminal and lateral buds will grow through the mesh or become entangled. Terminal buds outside the tube are vulnerable to herbivores. Tree deformities may develop if terminal buds catch on the sides of a tube. Protruding leaves and lateral branches are likely to be damaged.




Figure 10—Seedling enclosed within a solid barrier.

Preferred tube size will depend on the species and age of the tree being protected, as well as the predatory species. Short tubes may deter pocket gophers, but tubes 5 to 6 feet long are normally recommended to prevent deer and elk damage. The extended length allows terminal growth beyond normal deer and elk browse height. Tube diameter needs to be adequate to permit tree growth. Small tubes, however, may provide better protection from burrowing animals and generally cost less. Tubes 2 to 3 inches in diameter are generally recommended for Douglas fir and western larch, while 4- to 6-inch diameter tubes are generally recommended for pine and hardwood seedlings.

Some type of support or anchor is usually attached to keep tubes upright. Supports come in all sizes and shapes and are constructed from a variety of materials (e.g., plastic, wood, metal). Spiral stakes anchor more firmly in the ground than do straight stakes. Stakes with hooks are more readily fastened to the tubes. Some tubes come with attached ties or wires that can be easily fastened to supports. Support

mechanisms can often be purchased along with tubes. Again, the necessary support will depend on objectives and the animals in the area. Additional support for tubes to protect seedlings from gnawing pocket gophers is minimal because the tubes are usually short and buried a few inches in the soil. Tubes stationed in areas with ungulates, however, require substantial support. Ungulates knock tubes over when they rub against them and may pull the tube and seedling out of the ground. Metal fence poles are effective but expensive.

**Operation:** Tubes can be installed around seedlings either before or after planting. When installed prior to planting, a flexible plastic tube that closely matches the seedling height is wrapped around each seedling. Seedlings are then planted with a few inches of the tube below the soil surface to help anchor the tube. Difficulties experienced with this method include getting the tree planted in an upright position and at the proper depth. Other difficulties include firmly packing soil around the roots.

Tubes applied after planting are worked down over the seedling and then



Figure 11—Plastic sheets with slots that fasten together to hold the barrier in place once it is wrapped around a seedling. Barriers are supported with 1/2 x 1/2 inch wooden stakes.

anchors or supports are attached. Tubes split down the side can be opened for installation. The sides are secured once the tube is in place. Mesh tubes must be installed vertically with the leaders centered to reduce chances of the seedlings' terminal buds becoming entangled in the mesh. Supports should be placed immediately next to the tube.

**Advantages:** Properly installed tubes protect seedlings from most wildlife species. Tubes are not hazardous to humans or wildlife and they do not restrict wildlife access to other forage. Also, the reduced competition and the altered microclimate produced by some tubes may enhance seedling growth.

**Disadvantages:** Tubes are expensive and labor intensive to install. Large, solid-wall tubes require substantial space for storage and transport. Improperly chosen tubes, or tubes poorly installed, can cause seedling deformities or increase seedling mortality. The microclimate produced by some tubes may reduce seedling growth. Also, the high humidity in some tubes may increase problems with foliage diseases.



#### **Sources:**

ADPI Enterprises, Inc.  
American Forestry Technology, Inc.  
Davlyn Manufacturing Company, Inc.  
Design Packaging  
Forest Protection Products  
Forestry Suppliers, Inc.  
Naltex Plastics, Inc.  
Norplex, Inc.  
Orchard Supply Company  
Pacific Western Container  
The Tensar Corporation  
Treesentials Company

### **Conical Protectors**

**Function:** Conical protectors reduce bird and rodent access to newly planted seeds and young seedlings.

**Description:** Conical protectors are cone-shape wire enclosures constructed from #3 mesh hardware cloth. A parallelogram pattern (7-1/4 x 8 inches; interior angles of 60° and 120°) will make a cone with a base diameter of 4-1/2 inches. Approximately 750 screens can be cut to this size from a single roll (3 x 100 ft) of hardware cloth. Larger sizes can be made for taller seedlings.

**Operation:** Previously cut parallelograms are rolled into a cone and fastened by entwining the cut wires together. The base of the protector is secured by inserting it a couple of inches into the soil. Small cones are removed or replaced with larger cones as the tree requires additional space.

**Advantages:** Conical protectors are relatively inexpensive and reusable. Limited experience is required to construct and properly install conical protectors.

**Disadvantages:** Conical protectors need to be removed as the seedlings grow or they may interfere with growth and cause deformities.

### **Bud Caps**

**Function:** Bud caps are placed over the terminal bud to prevent browsing by ungulates.

**Description:** Bud caps can be constructed from any lightweight material that will fold, such as waterproof paper, fine mesh netting, or short pieces of mesh tubing. Fold the selected material to form a pocket that is slipped over the terminal bud. The cap is then fastened together with staples.

**Operation:** Bud caps are slipped over the terminal buds and held together with staples. Center the terminal bud within the cap. As a precaution do not cover more than 25% of a seedling's foliage and make sure that the stem is of an adequate size to support the cap.

**Advantages:** Bud caps are relatively inexpensive and require minimum skill to apply.

**Disadvantages:** Bud caps can cause heat damage to terminal buds. Bud caps provide no protection from rodents and may be pulled off the seedling by deer and elk.

### **Obstructions**

**Function:** Obstructions are used to inhibit access to trees by foraging wildlife.

**Description:** Obstructions can be anything that will serve as a barrier, such as tree limbs or forest debris.

**Operation:** Cover the seedling with debris, such that the debris does not interfere with tree growth but hinders access by animals. Seedlings may also be planted directly into existing debris.

**Advantages:** Implementation costs are limited to the expense of labor to construct obstructions.

**Disadvantages:** Obstructions provide protective cover for small mammals and may increase damage by rodents.

### **Squirrel Guards**

**Function:** Squirrel guards discourage tree squirrels from climbing trees.

**Description:** A squirrel guard is a 2-foot collar of metal or plastic that encircles the base of a tree. The guard is fastened together with wire attached to springs to allow for tree growth.

**Operation:** The guard is fastened around the base of a tree approximately 6 feet above the ground.

**Advantages:** Tree guards are easy to install and work well to protect isolated trees.

**Disadvantages:** Unless squirrel guards are placed on all trees, they are generally ineffective in forests or wherever tree branches normally become entwined.

## **Exclosures**

### **Fencing**

**Function:** Fences exclude wildlife from the protected resources. Drift fences are also used to alter wildlife movements or direct them to other damage reduction methods, such as traps.

**Description:** A vast array of designs and materials to construct fences exist. When selecting a fence design, consideration needs to be given to the overall objectives, soil types, vegetation, topography, maintenance requirements, availability and cost of materials, expected life span, labor, and visual impact. This section is not intended to serve as a comprehensive discussion on fence materials, design, or construction. For more detailed information refer to additional readings listed in the bibliography section. Fence designs and specifications are also available through the sources listed below.

Fences are commonly constructed of wire stretched between posts (metal or wood). The wire can be smooth, or barbed to restrict passage (Figure 12), or electrified to deliver a shock (Figure 13). Woven fences come in a variety of shapes and sizes. Historically, steel wire has been used to construct most fences. Plastics, however, are becoming more common. Wooden or metal rails also work, but are more expensive options.

Several designs for deer (Figure 14) and elk fencing have demonstrated some efficacy to prevent animal entry. Fences to keep out elk and deer should be a minimum of 8 feet and preferably 10 feet tall. Woven wires are much more effective at deterring ungulates than are strands of smooth or barbed wire. A combination of woven wire with strand wire installed immediately above it to provide additional height can be

effective. An electrified fence provides much better protection than a similar nonelectrified fence. Building a double or slanted fence adds depth, making the fence more difficult for ungulates to jump over. Flagging or other visible materials should be attached to make wire fences visible to prevent animals from running into them.

Woven wire (Figure 15) or solid fences are necessary to restrict rodent movements. The effective size of weave is dependent on the species of rodent. The bottom of the fence should be buried or fastened tightly to the ground to stop nonburrowing rodents. For burrowing rodents, such as pocket gophers and mountain beavers, the fence needs to be buried at least 3 feet below the soil surface. At the bottom the wire should be bent outward and upward, like a "J." The cup of the "J" needs to be approximately 6 inches wide with at least a 3-inch lip. The "J" is implemented because burrowing animals may dig downward along the outside of the buried fence. When they encounter the "J," it prevents them from continuing down and under the fence. An electrical wire or a slick sheet of metal (12 inches) fastened along the top will prevent most rodents from climbing over a fence.

**Operation:** Steps to construct a fence will depend on the selected design. However, most operations will include the following steps:

1. Coordinate objectives and planning with all appropriate persons
2. Gather site information and select a fence design
3. Secure easements if necessary
4. Arrange for materials and labor
5. Locate fence line
6. Identify locations for posts, braces, and gates
7. Set corner, gate, and line braces
8. Set line posts
9. Attach insulators for an electric fence
10. Stretch wire
11. Attach markers to make wires more visible
12. Ground wire fences
13. Set gates
14. Set and attach the energizer for an electric fence.




Figure 12—Strand of barbed wire spliced with the "Bull Wire Splice."




Figure 13—Closeup of "Super Wide Hot Tape" that can be strung between posts and electrified to protect agricultural products.

Fences need to be checked regularly to determine if repairs are necessary. Broken wires and posts need to be replaced or mended. Burrows and holes dug by rodents along the fence will need to be collapsed and filled. Vegetation should be mowed or clipped along electric fence installations to keep weeds from grounding the fence.

**Advantages:** Once constructed, enclosures protect resources for a long time and can be very effective. Fences are generally not dangerous to the physical well-being of wildlife or humans.

**Disadvantages:** Construction and maintenance of fences are expensive and labor intensive. Fences exclude wildlife from habitat and may interfere with normal activity and migration patterns. Wildlife can injure themselves if they run into or become entangled in fences.

**Sources:**

ADPI Enterprises, Inc.  
 American Feed & Farm  
 Baygard-Bay Mills, Ltd.  
 Brookside Industries, Inc.  
 Conwed Plastics  
 Dairyland Power Fence Company  
 Gallagher Power Fence, Inc.  
 Geotek, Inc.  
 Grassland Supply  
 Keystone Steel & Wire  
 K Fence Systems  
 Kiwi Fence  
 Live Wire Products, Inc.  
 Margo Supplies, Ltd.  
 Mississippi Valley Forest Products, Inc.  
 Naltex Plastics, Inc.  
 Premier Fence Systems  
 Qual-Line Fence Corporation  
 Safe Shop Tools  
 Southwest Power Fences  
 Techfence-Advanced Farm Systems  
 Tenax Corporation  
 Twin Mountain Supply Company  
 United Textile Corporation  
 Waterford Corporation  
 West Virginia Fence Corporation  
 Wildlife Control Technology, Inc.




Figure 14—Wooden fence installed to inhibit deer access to seedlings.




Figure 15—Rodent barrier constructed with metal aboveground and woven wire to extend 3 feet belowground when installed. Wire at the bottom will be bent to form a "J."

## Netting

**Function:** Netting is used to construct temporary exclosures, or where conditions require less weight or more flexible materials than generally used for conventional fences (Figure 16).

**Description:** Netting comes in a variety of weave sizes and patterns. Netting is flexible; it easily bends around corners and conforms to surface contours. Plastic and nylon are the most common materials. Again, the most appropriate style and size of netting depends on the objectives and available resources. Literature available through either the sources listed below or in the bibliography section provides alternative designs and product specifications.

**Operation:** Procedures to install netting are similar to those described for fencing. Net fences, however, are lightweight

and are generally temporary. Supports need not be as durable or as strong as those used for conventional fences. Netting can be hung from metal fence poles to create a barrier for deer or elk. A wire or bar strung across the top will secure the upper portion of the net and prevent sagging. Another wire stretched along the ground or stakes can be used to secure the bottom of the net.

Ungulate and bird depredation of seed beds can be inhibited by hanging nets over supports to create tent-like structures. Poles can be used to support netting. A guide wire stretched along the top of the poles will prevent sags between poles and permits poles to be spaced further apart. A series of inverted U's constructed out of plastic pipe also work well to support nets. Drape the net over the supports and then pull it outward and stake the edges securely to the ground.

**Advantages:** Netting is less expensive and easier to install than conventional fencing. Netting can be easily removed once conditions or objectives change. Netting is flexible and will easily cover irregular surfaces or bend around corners.

**Disadvantages:** Netting is more easily torn and needs repairs more frequently than do conventional fences. Like fences, netting excludes wildlife from habitat and may interfere with normal migration patterns. Animals can become entangled in nets and injure themselves.

### Sources:

ADPI Enterprises, Inc.  
APGAR, Inc.  
Birdbusters  
Bird-X, Inc.  
Blue Mountain Industries  
C. Frensch, Ltd.  
Conwed Plastics  
E.I. DuPont DeNemours and Co., Inc.  
Fuhrman Diversified, Inc.  
Green Valley Blueberry Farm  
Internet, Inc.  
J.A. Cissel Company, Inc.  
J.T. Eaton & Company, Inc.  
Laird Plastics, Inc.  
Margo Supplies, Ltd.  
Miller Net & Twine  
National Netting, Inc.  
Nichols Net & Twine Company, Inc.  
Nylon Net Company  
Orchard Equipment & Supply Company  
Prosoco, Inc.  
Sinco, Inc.  
Smith & Hawken  
Specialty Ag Equipment  
Sutton Ag Enterprises, Inc.  
Tenax Corporation  
United Textile Corporation  
Wildlife Control Technology, Inc.



Figure 16—Netting over support posts to inhibit entry by wildlife species.

# Traps

**T**rapping can be an effective means to remove animals that are damaging forest resources. Some trapping programs are selective in capturing specific individuals causing problems. Other trapping programs are implemented to reduce population densities of species that hinder reforestation efforts. These programs may be initiated after a problem is identified or in anticipation of problems on sites with high densities of species recognized as pests to forest resources.

The objectives of a trapping program will dictate the type and size of traps that should be utilized. In developing a trapping program several factors need to be considered, including the behavioral and biological characteristics of the target animal, ease of access to the trap site, experience and skill of the trapper, nontarget animals in the vicinity, cost effectiveness, state and federal laws and regulations, as well as other specific site considerations. Trap and release programs can be effective when specific individuals need to be removed. However, release sites need to be identified prior to capture. Euthanasia procedures and equipment also need to be ready before a program is implemented.

State and Federal laws and regulations govern the treatment of wildlife. Prior to implementing a trapping program, check with the proper authorities to ensure that your program is legal. Generally, a State's Department of Wildlife is an appropriate place to contact for information and guidance on trapping regulations.

Most traps pose a minimum of danger to humans. However, caution should be exercised, particularly when children may encounter trap sets. Additionally, trapped animals are usually agitated. Depending on their size, they can be very dangerous. Trappers need to exercise caution to avoid injuries inflicted while setting some types of traps (e.g., body-grip traps) and while handling captured animals.

## Live Traps

### Box Traps

**Function:** Box traps are used for the live capture of small- and medium-sized mammals, such as squirrels or snowshoe hares.

**Description:** Box traps (Figures 17, 18, and 19) are generally rectangular or square, with swing door(s) hinged at one or both ends of the trap. Most box traps have a treadle located near the center of the trap floor that releases the swing door(s) when depressed. Box traps are commonly manufactured of welded wire mesh. Traps constructed of plastic, wood, and steel are also commercially available. Solid wall traps (Figure 18) are applicable where exposure to extreme weather conditions, predators, or curious humans is a concern. Appropriate trap size depends on the species of concern (Table 1).

**Operation:** Box traps are used for the live capture of most small- and

medium-sized mammals. These traps, however, have demonstrated limited efficacy for the capture of canines and felines. Because of their bulk, box traps are difficult to transport. The triggering mechanisms may restrict their use to level ground. Capture success is greatly enhanced if traps are placed at the entrance of burrows or dens, adjacent to or within burrows or runways, or in close proximity of food sources. Bait is used to entice the target animal into the trap. Stress to captured animals can be reduced by checking the traps frequently, covering the trap with plastic sheeting or burlap, and ensuring that food and bedding material is available within the trap to maintain the animal until it is released.

**Advantages:** Box traps are used to capture animals alive and unharmed. Subsequently, the operator can release, relocate, or euthanize the animal, depending on the objective of the operation. Therefore, these traps pose little threat to nontarget species. Durability of these traps varies. However, most traps generally withstand repeated use if they are well maintained.




Figure 17—Several styles of box traps used to capture mammals.




Figure 18—Box trap with solid metal sides used to capture medium-sized mammals.




Figure 19—Modified Havahart trap specifically designed for the live capture of mountain beaver.

**Disadvantages:** Box traps are relatively expensive to purchase. Their bulk and weight make them difficult to transport. Trapped animals may injure themselves while attempting to escape or may undergo stress while in the trap. The size of traps hinders their versatility and does not readily permit them to be set on debris or uneven terrain.

**Sources:**

Animal Care Equipment & Services, Inc.  
 Animal Management, Inc.  
 B&G Chemicals & Equipment Co., Inc.  
 Bigelow Trap Company  
 B-Kind Animal Control Equipment  
 B.N. Bolton, Inc.  
 "Coon Getter" Traps and Animal Lures  
 Critter Control  
 Critter Gitters  
 Don Sprague Sales, Inc.  
 Duke Company  
 Ferrell and Company  
 Forestry Suppliers, Inc.  
 Goodart's  
 H.B. Sherman Traps, Inc.  
 J/J Fur Shop  
 Kemi-K Products, Inc.  
 Ketch-All Company  
 Krofick Outdoor Supply  
 Manufacturing Systems, Inc.  
 Minnesota Plasti-Catch  
 M&M Fur Company  
 Morrison Manufacturing Corporation  
 National Live Trap Corporation  
 Pioneer Wildlife Traps  
 R-P Outdoors  
 Safeguard Products, Inc.  
 Safety Live Trap  
 Seabright Enterprises, Ltd.  
 Stendal Products Inc.  
 Sterling Fur and Tool Company  
 Sullivan's Sure Catch Traps  
 Tomahawk Live Trap Company  
 Wickenkamp Live Trap Manufacturing  
 Woodstream Corporation

Table 1—Cage trap sizes that have been recommended for use to capture a selected list of representative mammals.

Mammal	Trap Dimensions	Mammal	Trap Dimensions
Ground squirrels	16" x 5" x 5"	Rabbits	32" x 9" x 9"
Rats	16" x 5" x 5"	Skunks	32" x 9" x 9"
Squirrels	19" x 6" x 6"	Opossums	32" x 10" x 12"
Squirrels	20" x 7" x 7"	Porcupines	40" x 12" x 12"
Muskrats	20" x 7" x 7"	Jackrabbits	42" x 15" x 15"
Mountain beavers	20" x 7" x 7"		

## Suitcase or Clamshell Trap

**Function:** Suitcase- (Bailey) or clamshell- (Hancock) type traps are most frequently used for the live capture of stream beavers.

**Description:** These traps consist of two metal frames, connected together

at one side with coil spring hinges. The frames are covered with chain-link wire mesh. When set, they resemble an open or half-open suitcase (Figure 20) or clamshell. A lever extends from the coil springs into the center of the trap and triggers the release mechanism. Only one size is available (21 x 39 inches closed) and there are no factory options or modifications. Individual traps weigh between 20 and 30 pounds.

**Operation:** These traps are generally used for the live capture of stream beaver. However, these traps can also be used to trap other aquatic mammals (e.g., river otter). Traps are generally set in or above water level near a lodge opening, adjacent to a slide along the shoreline, or on a frequently used

runway. Placement of beaver traps near lodges may be regulated by law in some states. Captured beavers may drown if the trap is set such that a portion of the trap does not remain above the water surface. Baits may be used to entice the target animal into the trap. Commonly used baits include corn, poplar, willow or alder branches, and castor lures. Traps should be checked at regular intervals to reduce the risk of captured animals drowning or becoming hypothermic, and to allow the release of nontarget captures.

**Advantages:** Properly set and baited suitcase or clamshell traps are highly selective for live trapping stream beaver. Further, any nontarget animals that are captured can be safely released.

These traps are safe and pose little danger of injury to stream beavers, nontarget species, or humans when set properly and checked frequently.

**Disadvantages:** Suitcase and clamshell traps are expensive and their bulk and weight makes them difficult to transport. Trapping with these traps is labor intensive. Their size restricts their use to areas that are accessible by vehicle or boat.

#### Sources:

Critter Control Wildlife Mgmt. Supplies  
Hancock Trap Company  
M&M Fur Company  
R-P Outdoors

## Leg-Hold Traps

**Function:** Leg-hold traps restrain the leg of an animal and are used to capture a broad range of species.

**Description:** Leg-hold traps are small, spring-activated mechanical capture devices. They consist of a pair of jaws, one or more springs, a triggering mechanism, and a circular base onto which the jaws and springs are attached. When pressure is applied to the triggering device, the jaws spring shut, capturing the leg of an animal. All leg-hold traps are manufactured with steel or steel alloys and are available in many sizes (Table 2). Three basic types of leg-hold traps are commercially available:

1. Coil spring
2. Under spring, or jump trap
3. Long spring.

Coil-spring traps (Figure 21) are powered by two or more small springs built into the center of the trap. The integral design of the coil-spring trap reduces its size and profile, permitting better concealment. Under-spring traps are powered by a single, integral spring. Long-spring traps (Figure 22) are powered by one or two large external "hinge" springs. These external springs increase the size and weight of the




Figure 20—Suitcase traps used to capture beaver.



long-spring trap. Leg-hold traps have been frequently modified or reconfigured to improve the perceived humaneness of the trap or to enhance its ability to catch and hold an animal. Common modifications include offset jaws, pan tension devices to exclude

smaller animals, and various swivels or mounting devices for securing the trap, and altering the spring strength or padding the jaws.

**Operation:** Leg-hold traps are commonly used to capture a wide variety of mammals. Disturbance by nontarget species can limit the use of these traps in some areas. Several factors need to be considered for proper trap sets, including but not limited to: behavior of the target species, habitat conditions, and the presence of nontarget species. In terrestrial settings, unbaited sets may be placed in or near travelways frequented by target animals. Baits or lures may be used to attract the target animal to a trap set. Aquatic sets are most commonly made in shallow water

near runways or adjacent to territorial marking sites used by the target animal. In general, coil-spring traps are used for smaller mammals in terrestrial settings, while long-spring traps are more commonly used in aquatic sets or for larger terrestrial animals.

**Advantages:** Leg-hold traps are significantly less expensive than cage traps. They cost substantially more than snares, however. Relative to cage traps, leg-hold traps are lightweight and small. Operators can transport 15 or more leg-hold traps, compared to one or two cage traps. These traps are generally durable, and replacement parts are readily available. A skilled trapper can place sets to minimize the incidental capture of nontarget animals. Generally, leg-hold traps permit the live release of nontarget animals. Human safety hazards associated with the use of leg-hold traps are minor.

**Disadvantages:** Leg-hold traps can inflict leg injuries to both target and nontarget animals. Injuries may result from the force of the trap or through efforts of the captured animal to escape. These injuries can be reduced by using offset or padded trap jaws, and by frequent checks of trap sets. Some expertise is required to effectively place trap sets and to minimize the risk of nontarget captures. The use of leg-hold traps is sometimes controversial and is banned in some states. Experience or training is required for leg-hold traps to be used safely and effectively.

#### Sources:

Bigelow Trap Company  
Duke Company  
Ferrell and Company  
Goodart's  
Leggett's Supplies  
Manufacturing Systems, Inc.  
Minnesota Trapline Products  
M&M Fur Company  
Montgomery Traps, Inc.  
O'Gorman Enterprises  
Ranchers Supply, Inc.  
R-P Outdoors  
Sterling Fur and Tool Company  
Woodstream Corporation

Table 2—Leg-hold trap sizes and jaw spreads that have been recommended to capture a selected list of representative mammals.

Mammal	Trap Size	Jaw Spread
Rats	#0	3.5"
Pocket gopher	#0	3.5"
Opossums	#1-#2	4" to 5.5"
Woodchucks	#1.5	4.75"
Skunks	#1.5	4.75"
Beaver	#4	5.5"




Figure 21—Coil spring leg-hold trap for the live capture of mammals.




Figure 22—Long spring leg-hold trap for the live capture of mammals.

## Foot Snares

**Function:** Foot snares can be used to capture a wide variety of animals. They are the most common and effective means used to capture bears, particularly in remote locations.

**Description:** A foot snare generally consists of a wire lead that is looped through a locking device and attached to a swivel (Figure 23). Some foot snares may be powered by a device that throws the snare upward onto the animal's leg. The snare is generally fixed to a tree or other stationary object, or to a drag to minimize an animal's movement away from the capture site. The type and diameter of cable, cable length, and type of lock device depends on the target animal (Table 3). Nontarget injuries and captures can be limited by placing a stop on the lead to



prevent closure beyond that necessary to hold the species of concern, or by using a break-away lock or cable that releases when force is exerted beyond that expected from the target species.

**Operation:** Foot snares are used to capture a wide variety of animals. They are the most common and effective means used to capture bears, particularly in remote locations. Snares can be placed in or near pathways frequently traveled by the target animal, or used in conjunction with baits or lures to entice the target animal to the snare. Considerable skill and knowledge of

animal behavior is required to make effective sets.

**Advantages:** Snares are the least expensive type of capture system. Their compact size and light weight make them easy to transport. Most snares are durable. Broken snares can readily be repaired with inexpensive replacement parts. Snares pose minimal safety hazards to humans and nontarget species.

**Disadvantages:** Considerable skill is required to place effective sets. Though snares themselves pose minimal

hazards to humans, captured animals have room to move and can inflict serious injury to intruders if approached too closely. This can be particularly dangerous if an animal has moved a drag set so the location of the animal is unknown. The use of snares may be regulated or restricted by law in some states.

#### Sources:

Grawes Animal Lures  
Hoffman Trapping  
Supplies, Ltd.  
M&M Fur Company  
Rocky Mountain Wildlife  
Products  
R-P Outdoors  
The Snare Shop

**Description:** Body-grip traps consist of two rectangular wire jaws with strong coil springs on one or both sides of the trap (Figure 24). The trap is armed by spreading the jaws apart against the force of the springs, which are then held in place with a notched lever or dog. Attached to the dog is a wire trigger that extends to the center section of the set trap. When the trigger is moved, it releases the dog, allowing the trap jaws to rapidly spring shut. Animals are killed with a strong blow to the body, often breaking the neck or spine. Body-grip traps are manufactured by several companies, all of which make traps in a number of standard-sized models (Table 4).

**Operation:** Body-grip traps can be used for the lethal removal of most small- to medium-sized mammals. Accidental encounters are generally lethal or injurious. Therefore, the successful release of nontarget species is unlikely. Caution needs to be exercised to ensure that nontarget exposures are minimized. Use of body-grip traps may be restricted by law in some states.

Trap placement is similar to that used for live traps. In terrestrial settings, unbaited sets are placed in or near travelways frequented by target animals, or baits or lures may be used to attract an animal to a trap set. These traps are often placed in burrows to remove fossorial rodents. In aquatic settings, sets are most commonly made in shallow water near runways or adjacent to territorial marking sites used by the target animals.

**Advantages:** Body-grip traps are relatively inexpensive. Their cost is similar to or less than the cost of leg-hold traps. Their light weight and compact construction makes body grip traps easy to transport. They are generally durable. A skilled trapper can use body-grip traps to make selective captures, especially for fossorial and aquatic species. Most animals are generally killed in a quick and humane manner.

## Lethal Traps

### Body-Grip Traps

**Function:** Body-grip traps are designed for the lethal removal of animals. Death is usually quick, generally induced by breaking the neck or the spine of the captured animal.




Figure 23—Foot snare attached to a spring; a typical foot snare set is depicted below.

Table 3—Snare sizes and cable sizes that have been recommended to capture a selected list of representative mammals.

Mammal	Snare Size	Cable Size
Squirrels	00-20"	1/32"
Rabbits	0S-30"	1/16"
Beavers	2S-40" to 2XX-72"	3/32"
Bears	5XX-120" to 6XX-120"	5/32" to 3/16"




Figure 24—Body-grip trap, and a cutaway of the trap set in a mountain beaver burrow.

Table 4—Body-grip trap sizes and jaw spreads that have been recommended to capture a selected list of representative mammals.

Mammal	Trap Size	Jaw Spread
Squirrels	110, 120	4-1/2" x 4-1/2"
Mountain beavers	110, 120	4-1/2" x 4-1/2"
Rabbits	120	4-1/2" x 4-1/2"
Skunks	120, 160, 220	4-1/2" x 4-1/2", 6" x 6", 7" x 7"
Opossums	120, 160, 220	4-1/2" x 4-1/2", 6" x 6", 7" x 7"
Porcupines	220, 330	7" x 7", 10" x 10"
Beavers	220, 280, 330	7" x 7", 8" x 8", 10" x 10"

**Disadvantages:** Body-grip traps do not allow for the selective release of nontarget animals. Therefore, caution must be exercised to avoid the capture of nontarget species. If a trap is improperly set, or is sprung by a nontarget animal, the trapped animal may sustain sublethal injuries and will need to be euthanized by the trapper.

voles; the larger, heavier (rat) traps can be used to capture chipmunks and squirrels, as well as rats. The Museum Special trap was designed by small mammal collectors to prevent skull damage and is effective for the capture of intermediate-sized rodents.

Body-grip traps pose substantial risk of injury for an untrained operator.

#### Sources:

Bigelow Trap Company  
Blue Mountain Industries  
Critter Control  
Duke Company  
Mitlying Development  
M&M Fur Company  
Pioneer Wildlife Traps  
R-P Outdoors  
Woodstream Corporation

## Snap Traps

**Function:** Snap traps are used for the lethal removal of small mammals.

**Description:** Snap traps (generally used for rats or mice) consist of a wire frame jaw attached to a wooden or plastic base, and powered with one or two coil springs (Figure 25). The trap is set by opening the jaw and securing it with a wire hook attached to a trigger. Movement of the trigger releases the jaw, pinning the animal between the jaw and the plate. Snap traps are commonly available in two sizes. The small (mouse) size is suitable for the capture of mice, shrews and

**Operation:** Snap traps are most commonly used to control commensal rodents in and around residential, agricultural, and industrial buildings. However, they are also used to capture small mammals in field situations. Traps are generally set on the ground or on a flat surface. Animals are enticed to the trap by bait placed on the trigger. Peanut butter mixed with oats, apple, raisins, or bacon is an effective bait for most rodents. Larger snap traps can be mounted vertically on a tree (pan downward) to capture squirrels. Traps used in damp environments must be treated with lacquer or wood sealer to avoid rapid deterioration. These traps also require careful cleaning and maintenance after each use. Risk of nontarget capture of birds may be reduced by enclosing traps within a shelter.

**Advantages:** Snap traps are the least expensive type of trap. They are lightweight and are easily transported. Most target animals are killed in a quick and humane manner. Human safety risks are minimal.

**Disadvantages:** Snap traps do not allow selective release of nontarget animals. Therefore, caution must be exercised in their use to avoid the capture of nontarget species. A captured animal may sustain sublethal injuries and will need to be euthanized. Snap traps are not very durable and wear out with repeated use.




Figure 25—A snap trap used for small rodents.

#### **Sources:**

B&G Chemicals & Equipment Co., Inc.  
Critter Control  
GMK Company  
Gremar Company  
Guardian Trap Company  
Hadley Products Company  
Ketch-All Company  
Kness Manufacturing Company, Inc.  
McGill Metal Products Company  
Micro-Gen Equipment Corporation  
M&M Fur Company  
Pest Control Supplies  
P-W Manufacturing  
Rodent Control Systems, Inc.  
R-P Outdoors  
Woodstream Corporation

### **Pocket Gopher Traps**

**Function:** Pocket gopher traps are specifically designed for the lethal removal of pocket gophers.

**Description:** Two general types of pocket gopher traps are commercially available, the box choker (Figure 26) and pincher traps (Figures 27 and 28). The box choker trap is basically a snap trap enclosed within a box that has one end open. Bait may be used to attract a pocket gopher to the rear of the box. A wire frame jaw, released when the trigger located at the far end of the box is moved, pinions the animal to the base of the box. Pincher traps, such as the Macabee, have two jaws (or a wire loop and one jaw) that close together to pinion the pocket gopher. Jaws may be pointed inward to kill the animal more quickly.

**Operation:** Traps are placed within active pocket gopher burrows where fresh mounds or plugs are indicators of recent activity. Bait is generally not used with pincher trap sets. Slight openings to the surface behind a trap

set can entice a pocket gopher to the set when it attempts to plug the hole.

**Advantages:** Pocket gopher traps, relative to most other traps, are inexpensive. They are small and light, so an operator can readily carry a substantial number of them. Most are durable and can be used repeatedly. Nontarget animals are generally not a concern, since traps are set within active pocket gopher burrows. Pocket gophers are killed quickly and humanely. Risks to humans are minimal.

**Disadvantages:** Pincher traps do not permit the release of nontarget captures. Effective trap sets require training or an experienced operator. Pocket gopher trapping on large areas is labor intensive.

#### **Sources:**

Don Sprague Sales, Inc.  
F.B.N. Plastics  
P-W Manufacturing  
R-P Outdoors  
Wilco Distributors, Inc.  
Windear Manufacturing Company  
Z.A. Macabee Gopher Trap Company

### **Neck Snares**

**Function:** Neck snares (Figure 29) are used for the lethal capture of animals. Neck snares can also be used to capture animals alive if a stop is placed on the snare to prevent closure.

**Description:** Snares generally consist of a cable lead that is looped through a locking device and attached to a swivel. When an animal is captured, the loop tightens as pressure is applied against the cable. The cable diameter and length vary with the intended use and the target animal (Table 3). Nontarget

animal injuries can be limited by placing a stop on the lead to prevent closure beyond that necessary for the target animal, or through a breakaway system that releases if a force is exerted beyond that expected from the target animal.

**Operation:** Snares can be used to capture a variety of mammals. Sets are placed along paths frequented by the targeted animal or are used in conjunction with baits. Considerable skill and knowledge of animal behavior are required for effective sets and to limit nontarget encounters. Captured animals are asphyxiated. The loop tightens around the animal's neck and death generally occurs within minutes. Aquatic trap sets are generally placed such that the animal is held beneath the water.

**Advantages:** Snares are the least expensive type of trap. Their compact size and light weight make them easy to transport. Most snares are very durable. Broken snares can be readily repaired with inexpensive replacement parts. Human safety hazards are minimal.

**Disadvantages:** Neck snares may not permit the release of nontarget captures. Considerable skill is required to use neck snares effectively and to minimize risks to nontarget species. The use of these devices may be highly regulated or prohibited by law in some states.

#### **Sources:**

Blue Mountain Industries  
Grawes Animal Lures  
Hoffman Trapping Supplies, Ltd.  
M&M Fur Company  
Rocky Mountain Wildlife Products  
R-P Outdoors  
The Snare Shop  
Woodstream Corporation




Figure 26—A set pocket gopher box choker trap.




Figure 27—One style of pincher trap used for pocket gophers.




Figure 28—A pincher trap used for pocket gophers and a cutaway of a trap set in a pocket gopher burrow.




Figure 29—A neck snare and a typical neck snare set.

# Toxicants

**T**oxicants can be an effective means of quickly reducing high populations of some problem animals, or of maintaining acceptable population densities. Generally, they are most applicable where existing or potential damage problems are related to population density. However, they may also be appropriate for treating areas with limited access or where other approaches have proven ineffective.

Toxicants are generally classified within one of three categories depending on their mode of action. Acute toxicants are ingested and are most often lethal soon after a single encounter. Chronic toxicants are also ingested, but are not immediately lethal. Fumigants are lethal gases that are injected into the burrows of the targeted species. The most applicable approach will be directed by your overall objectives and any restrictions regulating toxicants that apply to your specific situation.

Toxicants pose a potential hazard to the operator and other humans, as well as to nontarget species. Therefore, extreme caution needs to be exercised in handling, storing, and applying toxicants, and in disposing of waste materials. The applicator assumes responsibility to ensure that all legal and safety concerns are met prior to using toxicants. Carefully read and follow the restrictions and safety precautions provided on labels for each product. Federal and state registrations certify that it is legal to use the product according to the conditions and restrictions stipulated on the approved label. Registration does not guarantee the availability or the efficacy of a product. Rodenticides are commonly used to protect forest resources from depredation by pocket gophers (*Thomomys* spp.) and to a lesser extent by mountain beaver (*Aplodontia rufa*), voles (*Microtus* spp.) and woodrats (*Neotoma* spp.). At present, there are no toxicants registered for application in forest habitat to control beaver (*Castor canadensis*), porcupines (*Erethizon dorsatum*), squirrels (*Sciurus* spp.) or rabbits (*Sylvilagus* spp., *Lepus* spp.).

Toxicants are probably not feasible treatments for most nonrodent species that inflict damage to forest resources.

## Ingested Toxicants

### Acute Toxicants

**Function:** Acute toxicants are used to quickly reduce populations of rodents.

**Description:** Single encounters with acute toxicants can be lethal to rodents. Death, depending on the toxicant, results from paralysis of the respiratory or central nervous system, cardiac failure, gastrointestinal irritation, hemolysis, destruction of kidney and liver tissue, or hypothermia.

Several acute toxicants are available, including strychnine and zinc phosphide. Strychnine induces paralysis of the central nervous system. Death occurs within 30 minutes if a lethal dose is ingested. The bitter taste of strychnine can hinder target animal acceptance of treated bait. Zinc phosphide produces phosphine gas when it reaches the stomach or intestines. The gas causes heart failure and liver or kidney damage. Symptoms generally occur within 30 minutes after intake and death occurs within 24 hours. Zinc phosphide has a distinctive garlic odor that may be attractive to some rodents at low concentrations.

**Operation:** Intake of lethal doses of toxicants by rodents can be problematic. Rodents may be hesitant to initially accept treated baits, or if sublethal doses are ingested, they may develop bait-shyness. Bait-shyness is a learned food aversion formed when a food is paired with gastrointestinal illness. Once bait-shyness occurs, the likelihood of subsequent intake is doubtful. Prebaiting can enhance the acceptance of treated baits. Before treatment, present untreated bait for target animal

consumption. Mimic all procedures (e.g., bait location) that will be used in the subsequent program. Once untreated baits are being readily ingested by the target species, switch to the toxic bait.

The type of bait and its placement are important to the efficacy of a program to reduce populations of target species, and to minimize potential risks to nontarget species. Foods that are highly preferred by the target species and that do not spoil rapidly make the best baits. Most toxicants are available in ready-to-use baits (Figure 30). While this may increase the cost, it also reduces chances for accidental poisoning and incorrectly prepared baits.

Baits need to be placed in burrows or other locations where they will be repeatedly encountered by the target species yet not be readily available for nontarget species. Toxicants must be applied according to the regulations and restrictions on the applicable labels.

**Advantages:** A quick response can be expected with an effective program using acute toxicants. When delays in reducing rodent populations will result in extensive resource losses, acute toxicants may be the only reasonable alternative. Generally, acute toxicants are a relatively inexpensive means to reduce rodent populations. Single treatments require small amounts of toxicants and little labor to apply.

**Disadvantages:** Sublethal doses may induce bait-shyness, rendering a population reduction program ineffective. Prebaiting may be necessary, which enhances the efficacy but also increases the cost of a program. Acute toxicants are lethal to a broad spectrum of animals. Single encounters with these products can accidentally poison nontarget species.

### Sources:

Archem Corporation  
Bell Laboratories, Inc.  
B&G Chemicals & Equipment Co., Inc.  
Boehringer Ingelheim Animal Health, Inc.  
Bonide Products, Inc.



Figure 30—Mole and gopher bait.

Chas. H. Lilly  
 C. J. Martin Company  
 Eckroat Seed Company  
 Fort Dodge Chemical Company  
 HACO, Inc.  
 LiphaTech, Inc.  
 Monterey Chemical Company  
 Motomco, Ltd.  
 Nevada State Department of Agriculture  
 Nott Manufacturing Company, Inc.  
 Petersen Seeds, Inc.  
 Pocatello Supply Depot  
 Purina Mills, Inc.  
 RCO, Inc.  
 Rue R. Elston, Inc.  
 Sebesta Bait Mixing Plant  
 South Dakota Department of Agriculture  
 Southwest Chemical Company  
 VanWaters and Rogers  
 Wilbur-Ellis Company  
 Wilco Distributors  
 York Distributors

## Chronic Toxicants

**Function:** Chronic toxicants are not immediately lethal to rodents, or lethal doses are generally ingested only after multiple encounters.

**Description:** Anticoagulants interfere with the production of vitamin K, which is essential for the synthesis of prothrombin and other factors

necessary for the natural clotting of blood. Inhibition of blood clotting, combined with the disruption of small blood vessels, causes fatal internal bleeding. The effects of most anticoagulants are cumulative. Lethal doses require repeated ingestion. Single-dose anticoagulants are not acute, since death occurs only after several days. However, like acute toxicants, they require only a single exposure. With single-dose anticoagulants, feeding stops

immediately. Death can occur after only 2 days. Though the mode of action for anticoagulants is relatively slow, they are reported to induce a painless death. Nontarget animals that have ingested anticoagulants can be treated with supplements of vitamin K, an effective antidote.

Anticoagulants belong to one of two groups, the hydroxycoumarins and the indandiones, which differ in chemical structure but are similar in their physiological effect. The hydroxycoumarin group includes brodifacoum and warfarin among others, while chlorophacinone and diphacinone belong to the indandione group.

Warfarin was the first anticoagulant developed. It is substantially more toxic when it is ingested over a period of consecutive days rather than during a single feeding bout. Bromadiolone was among a second generation of anticoagulants developed specifically to target warfarin-resistant rodents. Bromadiolone will induce death after a single feeding when high concentrations are ingested. Reduced concentrations are lethal after multiple feedings. Chlorophacinone and diphacinone are substantially more toxic than warfarin, but are not generally effective against warfarin-resistant rodents.

Cholecalciferol may be a single- or multiple-dose rodenticide. It causes calcium to mobilize from the bone matrix to plasma, which in turn induces hypercalcemia. After ingesting a lethal dose, an animal will quit eating. Death occurs within 3 to 4 days.

**Operation:** An effective program requires that multiple dose anticoagulants be ingested repeatedly over a period of time. Therefore, the bait should be readily acceptable and available for ingestion by the target species. Prepared baits with anticoagulants are commercially available. Bait stations ease repeated applications. Considerations for the operational application of chronic toxicants are similar to those for treatment with acute toxicants. Prebaiting, however, is not necessary because rodents generally do not develop bait-shyness to anticoagulants. Nontarget hazards are also less likely because single exposures are usually not lethal and antidotal treatment is possible.

**Advantages:** Chronic toxicants can be highly effective in reducing populations of target species. Hazards to nontarget species from a single encounter are slight and an antidote (vitamin K) is available. Anticoagulants are not likely to induce bait-shyness. Therefore, prebaiting is not necessary. The mode of action suggests an apparently painless death for the target species.

**Disadvantages:** Chronic toxicants may be effective only after being ingested on several subsequent encounters. Thus, populations of target species are not reduced as quickly with chronic toxicants as they would be with acute toxicants. Operational control with anticoagulants can also be expensive because of high labor input and the high cost of materials due to repeated baiting.

**Sources:**  
 Archem Corporation  
 Bell Laboratories, Inc.  
 B&G Chemicals & Equipment Co., Inc.  
 Boehringer Ingelheim Animal Health, Inc.

Bonide Products, Inc.  
 C.J. Martin Company  
 Ferrett Laboratories, Inc.  
 HACO, Inc.  
 J.C. Ehrlich Chemical Company  
 J.T. Eaton & Company, Inc.  
 LiphaTech, Inc.  
 Motomco, Ltd.  
 Nott Manufacturing Company, Inc.  
 Pest Control Supplies  
 Pocatello Supply Depot  
 Purina Mills, Inc.  
 RCO, Inc.  
 South Dakota Department of Agriculture  
 Southwest Chemical Company  
 VanWaters and Rogers  
 Wilbur Ellis Company  
 Wilco Distributors, Inc.  
 York Distributors

rodent burrows. Phosphine and carbon monoxide are two examples of fumigants. Aluminum phosphide reacts with atmospheric or soil moisture to release phosphine gas. Phosphine is colorless and produces only a slight odor, similar to that of carbide. Phosphine gas is dangerous at concentrations as low as 200 ppm (parts per million). At concentrations of 400 ppm, it is lethal to humans. At concentrations of 1000 ppm, a few breaths will kill humans.

Gas cartridges are incendiary devices that produce smoke and carbon monoxide when ignited. Concentrations of 200 ppm may induce symptoms of carbon monoxide poisoning within a few hours. Exposures to concentrations approaching 1000 ppm will induce death within 4 hours.

**Operation:** First, ensure that no nontarget species are present. Then check for and seal off all exits or porous areas where the gas may leak out. Smoke bombs are generally effective in locating numerous burrow openings. Once the systems are sealed, fumigants are inserted into the underground systems. Carefully read and follow instructions provided on the product labels.

Moisture content and soil particle size are important considerations for fumigation. Fumigants are most effective when the soil moisture is high and the soil temperature is greater than 60°F. Fumigation failures most frequently occur in dry, porous soils.

Extreme caution needs to be exercised when handling fumigants. They are lethal to all animals subjected to them. All restrictions and directions provided on the appropriate labels should be carefully followed.

**Advantages:** There are few advantages to using fumigants as a primary means to reduce populations of forest rodents. Fumigants may provide a feasible alternative when other measures are ineffective.

**Disadvantages:** Fumigants pose considerable hazards to nontarget animals that coexist in burrows with target species. They can also be extremely hazardous to the applicator. Operational use of fumigants is generally more expensive than other toxicants.

#### Sources:

Bernardo Chemicals, Ltd.  
 Degesch America, Inc.  
 Douglas Products and Packaging Co.  
 Nott Manufacturing Company, Inc.  
 Pestcon System, Inc.  
 Pocatello Supply Depot  
 Research Products Co.  
 Roxide International, Inc.  
 VanWaters and Rogers

## Fumigants

**Function:** Fumigants (Figure 31) are used to produce a toxic atmosphere within the burrows of rodents.

**Description:** Fumigants are poisonous gases or compounds that produce poisonous gases. They are injected into




Figure 31—Fumigant pellets.

## Applicator Equipment

### Broadcast Applicators

**Function:** Broadcast applicators dispense baits over a wide swath.

**Description:** Hand broadcast spreaders or mechanical broadcast spreaders are commercially available. Most broadcast seeding equipment, with minor modifications, can be used to broadcast seed or pellet baits. Hand broadcasters consist of a storage bin, a crank-operated distribution device and an adjustable metering slot. The adjustable slot governs the rate of application. Mechanical broadcasters are similar to the hand-held versions, but have motorized components.

**Operation:** After filling the bin with the desired baits and adjusting the application rate, the operator traverses the specified site. Baits are dispersed at even intervals across the soil surface.

**Advantages:** Baits can be rapidly spread across treated sites. Broadcast applicators are most effective when

applying baits for rodents that forage on the surface, such as mice and voles. They also minimize direct contact with baits by the operator.

**Disadvantages:** Spreading baits across the soil surface increases the chance for encounters by nontarget species. Most species detrimental to forest resources can be more specifically targeted by placing baits within their burrow systems or on runways. Greater amounts of toxicants may be needed than with other methods because encounters are by chance rather than with strategically placed baits.

**Sources:**

Davenport Seed Corporation  
Herd Seeder Company

## Burrow-Builders

**Function:** Burrow-builders construct artificial burrows and automatically dispense bait underground for pocket gophers.

**Description:** A tractor-drawn unit is available in hydraulically operated three-point hitch models. The burrow-builder consists of a coulter blade to cut roots of plants ahead of the knife, a knife and torpedo assembly to construct the artificial burrow, a bait bin with a metering device to dispense baits at the desired rate, a dispensing mechanism, and a packer wheel to close the burrow.

Similar machinery can be used to dispense bait for voles. Voles prefer to travel along trails and are unlikely to encounter most bait spread at random. Instead of dispensing bait in burrows, the machine dispenses bait along a path created along the ground surface.

**Operation:** The tractor is driven across the treated site, constructing burrows and automatically dispensing bait. Artificial burrows should be constructed at depths similar to those for the natural pocket gopher burrows found in the area and spaced at 20- to 25-foot

intervals. Burrows that encircle the perimeter of the treated site will help to inhibit pocket gopher invasions from areas outside the site. However, continuous burrows should not be extended across the treated site as these may be used by pocket gophers for rapid movement across the site.

While building burrows the operator should periodically check to ensure that the bait dispenser is not plugged and that the burrows are in good condition. Wet soil and debris may collect on the packer wheel or knife shank, creating inadequate burrows. Burrows created in extremely dry soil tend to collapse.

**Advantages:** Burrow-builders or trail-builders can rapidly and economically treat large acreages. Toxicants are delivered beneath the surface or on trails where they are encountered by pocket gophers or voles. This approach increases the likelihood of encounters by pocket gophers or voles, while decreasing the incidence of nontarget encounters.

**Disadvantages:** Burrow-builders or trail-builders are not applicable when treating small acreage, or treating sites with excessive debris or steep terrain.

**Sources:**

The Perry Company  
RCO, Inc.  
Rue R. Elston Company  
Western Alfalfa  
Wilco Distributors, Inc.

## Bait Stations

**Function:** Bait stations (Figure 32) permit a continuous supply of bait to be available to the target species, while hindering access to nontarget animals.

**Description:** A variety of bait stations made of plastic, cardboard, or

metal are available through commercial suppliers. They generally consist of a box with an opening suitable for entry by the target species and some kind of bar or lip to prevent the bait from being easily carted out of the container. Homemade bait stations can be easily created by using old tires, a pipe, or an enclosed box. The bait station should be clearly labeled with "poison" or similar warning, and secured to prevent it from being knocked or dragged out of place. Some rodenticide uses may require tamper-proof bait stations to reduce human or nontarget hazards. Check the pesticide labeling carefully before beginning a program.

**Operation:** Bait stations need to be placed in the field a week or so prior to treating with baits. Prebaiting with nontoxic foods may be beneficial. Once the target animals become accustomed to the station, bait is placed inside. Bait stations need to be checked frequently to replace ingested or contaminated bait. After the desired effect is achieved, bait stations need to be removed. Any remaining bait should be returned to its original container or disposed of properly.

**Advantages:** Bait stations are relatively inexpensive and easy to maintain. They minimize environmental hazards and risks to nontarget species and can be safely handled by humans. A continuous




Figure 32—Assorted bait stations for small rodents.



supply of bait can be maintained in good condition; this is particularly important for the use of anticoagulants, or when rapid invasion occurs on treated sites.

**Disadvantages:** Most species detrimental to forest resources can be more specifically targeted by placing baits within their burrow systems. Prolonged maintenance of bait stations can become expensive as they require more intensive labor than other delivery methods.

**Sources:**

B&G Chemicals and Equipment Co., Inc.  
Bell Laboratories, Inc.  
Eco-Bait  
Geyer Irrigation  
The Huge Company  
J. T. Eaton & Company, Inc.  
LiphaTech, Inc.  
Motomco, Ltd.  
Solvit, Inc.  
Wilco Distributors, Inc.  
Woodstream Corporation

## Probes and Bait Dispensers

**Function:** A probe can be used to locate burrows and create an opening to drop bait. A bait dispenser is a probe

with an automatic bait dispenser attached.

**Description:** Probes can be made of almost any rounded material that can be used to penetrate the soil. A broom or shovel handle works quite well in soft soil for moderate use. A more durable probe works better when the soil becomes harder or for more extensive efforts. Probes generally look like a large "T" that is about 30 inches long with a handle 12 to 15 inches long across the top. A foot rest may also be attached to aid in penetrating hard surfaces. Bait dispensers are similar, except that a bait container is attached, along with a mechanism that dispenses measured doses of bait when activated.

**Operation:** The operator probes the soil around mounds where an active pocket gopher system is likely to be encountered. Fresh mounds or plugs are an indication of recent activity. An experienced operator can feel the probe strike a burrow by the reduced resistance as the probe enters the open space.

Once a system is located, the operator can dig out the system and insert bait with a spoon, or drop the bait through the opening created by the probe. A

hollow tube inserted through the probe hole helps to deliver bait to the burrow. The bait dispenser is operated in a similar manner. Once a burrow is located, the delivery mechanism is triggered and the bait is automatically dispensed into the burrow. Regardless of the approach, the entry hole needs to be completely sealed or the pocket gopher may plug the hole and cover the bait with soil while doing so.

**Advantages:** Probing, and then directly inserting bait, ensures that the bait is delivered into the burrow system. Exposure of baits to nontarget species is minimal with these methods.

**Disadvantages:** Hand baiting large areas is labor intensive. Skilled operators are necessary so that burrows can be readily located and the bait properly placed. New personnel will require substantial training.

**Sources:**

Don Sprague Sales, Inc.  
Eckroat Seed Company  
Fuhrman Diversified, Inc.  
Quinn Manufacturing Company  
Rue R. Elston, Inc.  
Southwest Chemical Company  
Topo Manufacturing and Welding Co.  
Wilco Distributors, Inc.

## Repellents (Chemosensory)

**W**ildlife damage to forest resources is predominately inflicted by foraging animals. Seedlings and sapwood are an attractive food resource for a variety of wildlife. The extent to which wildlife forage on these resources depends on the availability and relative desirability of alternative foods. Repellents (Figures 33 and 34) can be applied to seedlings to render them less attractive than the alternative foods. Consequently, animals select for plants or foraging areas other than those protected with repellents. The avoidance of repellents by wildlife may be innate or acquired through a conditioned food aversion.




Figure 33—"Big Game Repellent" used to inhibit foraging by herbivores.

Repellents that elicit initial avoidance are generally either irritants (e.g., capsaicin) or those that evoke a "fear" response (e.g., predator scents). These repellents require no prior conditioning to elicit avoidance behavior. Irritants stimulate trigeminal pain receptors. For example, jalapeño peppers contain capsaicin that activates pain receptors in mammals when ingested. Fear-inducing repellents are usually animal waste products or predator odors. Degrading waste products and most predator urine emit sulfurous odors.



Figure 34—"Tree Guard" repellent.

Conditioned food aversions occur when ingestion of novel foods is paired with nausea. Thus, any flavor paired with gastrointestinal distress can become an effective repellent. Efficacy of repellents based on conditioned aversions, however, are generally limited because animals must first be conditioned to avoid these materials. Further, the stimulus must be novel for the animals to form a strong aversion. Damage inflicted to seedlings during conditioning or subsequent sampling may be extensive. The use of repellents based on conditioned aversions can be especially problematic if the damage is inflicted by a transitory or a migratory species (i.e., deer moving from summer to winter ranges).

Wildlife responses to repellents vary among individuals, as well as among species. Obviously, responses to repellents that require conditioning will reflect each animal's prior experience. Less obvious, however, are response differences that occur because some

species are more responsive to certain stimuli than others. Sensitivity to "irritating" agents varies among species. Agents that stimulate trigeminal pain receptors in some species fail to stimulate similar receptors in other species. For example, mammals tend to avoid capsaicin, while birds are indifferent to it. Conversely, low concentrations of anthranilates are irritating to birds while similar concentrations are not offensive to mammals.

Response differences among wildlife also appear to reflect their foraging strategies. Carnivores (e.g., dogs) and omnivores (e.g., bears) avoid bittering agents, while most herbivores (e.g., deer, pocket gophers) are indifferent to them unless they have been previously paired with nausea. Likewise, meat eaters, such as coyotes and bears, are attracted to sulfurous odors, while most herbivores usually avoid substances with these odors.

A repellent's efficacy also depends on the desirability of the protected plant as a food source. Alternative forages are more likely to be selected if the protected plant is not a valued food source. Preferred plants (e.g., western red cedar) are more difficult to protect than plants that are not preferred (e.g., foxglove).

An effective program to reduce wildlife foraging using repellents also depends on the availability of alternative forage. An abundance of alternative forage permits animals to readily direct their destructive behavior toward other plants. After treatment, an animal's foraging choices will depend on the size of the protected area relative to its territorial boundaries. Species with vague or extensive territories, such as deer or bear, can more easily move to new areas to locate alternative forages than can species (e.g., pocket gophers) with small and more rigid territorial boundaries.

Foraging pressure on protected plants also depends on the types and densities of wildlife species. Competition among and within species may cause animals to select less preferred foods. Likewise, high population densities may limit foraging alternatives, rendering repellents less effective.

Some repellents, if improperly applied, can pose a human or environmental hazard. Therefore, caution must be exercised in the handling, storing, applying, and disposing of repellent materials. The applicator assumes responsibility for ensuring that all legal and safety precautions are followed. Restrictions and safety precautions provided on product labels must be carefully read and followed. Federal and State registrations certify that it is legal to use the product according to the conditions and restrictions stipulated on the approved label. Registration does not guarantee the availability or the efficacy of a product.

Generally, chemosensory repellents are categorized as those that deter animals from an area or reduce direct contact with a plant.

## Area Repellents

**Function:** Area repellents inhibit wildlife presence near the vulnerable forest resource.

**Description:** Area repellents emit odors that are undesirable to the target species. Degrading animal waste products and predator odors produce sulfurous odors that are generally offensive to most herbivores. A number of commercially available repellents incorporate these and other products into area repellents. Homemade remedies, such as bags filled with human hair or soap bars, have also been reported to reduce damage inflicted by foraging wildlife.

**Operation:** Place area repellents within the area to be protected following instructions on the product's label. The distance between repellent materials, and the amounts applied at each spot will depend on the product. Efficacy claims may report it to cover an extensive area or protect only within the immediate vicinity of the application. After the repellent is dispersed, monitor animal activity to determine whether it is working.

Several options need to be considered if the program is not effective. Additional material may be required if the original application was inadequate or weather conditions are inhibiting performance. Rain may dilute the repellent. Odors may be less volatile during cold temperatures. An alternative type of repellent may be more effective. Repellent products are not equally effective at deterring all species. Periodic switching of repellent types may decrease animal habituation. It may also be necessary to supplement or replace the repellent program with another approach to reduce damage.

**Advantages:** Area repellents protect all plants and plant parts within a treated area and are generally not harmful to wildlife or humans. Some area repellents can be an inexpensive means to protect forest resources and require little equipment or few hours of work to apply.

**Disadvantages:** Depending on the type of repellent and the number of applications, this approach can become expensive and labor intensive. Animals habituate to most repellents fairly quickly. Effective area repellents can potentially exclude wildlife from favorable habitat and other forage resources.

### Sources:

Avitrol Corporation  
Boehringer Ingelheim Animal Health, Inc.  
Bonide Products, Inc.  
Bushwacker Backpack and Supply  
Champon 100% Natural Products, Inc.  
Coughlan Products Corporation  
Degesh America, Inc.  
Dragon Corporation  
Dr. T's Nature Products Company, Inc.  
Faesy and Besthoff, Inc.  
Farnam Companies, Inc.  
Gustafson, Inc.  
HACO, Inc.  
IntAgra, Inc.  
Integral Designs  
J.T. Eaton & Company, Inc.  
Lakeshore Enterprises  
Margo Supplies, Ltd.  
Miller Chemical & Fertilizer Corporation  
Nature's All Natural Products, Inc.  
Nortech Forest Products, Inc.  
Pace International, Ltd.  
Plant Pro-Tech, Inc.  
PMC Specialties Group, Inc.

## Contact Repellents

**Function:** Contact repellents render plants less desirable for consumption by animals.

**Description:** Contact repellents are placed on or absorbed by the protected resource, and are generally avoided because of an aversive taste or smell. Animals enter areas treated with effective contact repellents, but do not ingest the treated plants. Most contact repellents are applied directly to the plant. Systemic repellents, however, are applied to the soil and absorbed by the plant. Many commercially available area repellents also work as contact repellents.

**Operation:** First, check to ensure that the repellent is not toxic to the plant. Treat and observe a few plants if phytotoxicity data are not provided on the label.

Repellents must be applied according to the restrictions and directions provided on the label. For the best results, treat each individual seedling. On large areas, however, it may be possible to treat only those seedlings located along the outside edge (100 feet) of the plot. Treating only the exterior seedlings reduces labor and material costs and may work if animals do not normally cross the untreated portion. Observe areas to determine whether wildlife are avoiding treated plants.

Ineffective efforts to reduce damage with repellents generally occur because an ineffective repellent was applied, or because the delay between treatments was too great. Few, if any, of the repellents currently available remain

active for more than 3 or 4 months. Weather conditions, particularly precipitation, may further decrease the period during which plants are protected by repellents.

**Advantages:** Contact repellents are generally not harmful to wildlife or to humans. These repellents protect specific plants and do not hinder wildlife use of other forage or habitat.

**Disadvantages:** Depending on the cost of the repellents and the need for repeated applications, this approach can be expensive and labor extensive. Animals tend to habituate to most repellents fairly quickly. Some repellents may be hazardous to humans if not handled carefully, or detrimental to the environment if not applied correctly. Extreme weather conditions may dilute contact repellents and make them ineffective.

**Sources:**

Animal Repellents, Inc.  
Boehringer Ingelheim Animal Health, Inc.  
Bonide Products, Inc.  
Burlington Scientific Corporation  
Bushwacker Backpack and Supply  
Champon 100% Natural Products, Inc.  
Coughlan Products Corporation  
Degesh America, Inc.  
Dr. T's Nature Products Company, Inc.  
Earl May Seed & Nursery Company  
Gustafson, Inc.  
HACO, Inc.  
IntAgra, Inc.  
Integral Designs  
J.C. Ehrlich Chemical Company  
Lakeshore Enterprises  
Margo Supplies, Ltd.  
Miller Chemical and Fertilizer Corp.  
Nature's All Natural Products, Inc.  
Nortech Forest Products, Inc.  
Nott Manufacturing Company, Inc.  
Pace International, Ltd.  
PMC Specialties Group, Inc.  
Sudbury Laboratories, Inc.  
Wilbur-Ellis Company

# Frightening Devices

**V**isual displays or noises that serve to alarm or appear threatening can be installed to frighten wildlife away from forest resources. Yelling at wildlife ingesting agricultural products was probably the first historical effort to prevent animal damage. Modern noisemakers are still used to scare animals away from resources. Visual displays, such as scarecrows, are also traditional means to alleviate depredation in gardens or grain fields. Modern displays are generally either more realistic in appearance or less cumbersome and easier to install than the traditional scarecrow. Frightening devices are most commonly used to deter birds, but they may also be used to deter mammals.

Animals are generally wary of any unfamiliar sound or sight, but they become less wary with time unless the noise or vision is paired with a negative reinforcer. Familiarity of wildlife to frightening devices can be minimized by installing or operating the devices only during periods when resources are most susceptible to damage. Devices, however, need to be used immediately after the onset of damage. Established movements and behaviors are much more difficult to disrupt than are newly forming behavioral patterns.

Efficacy of products can usually be increased by alternating techniques or use patterns. Sporadic displays or devices that are activated by an animal's presence are more effective than permanent or routine displays. Visual displays combined with noisemakers are generally more effective than either technique implemented alone. For example, sirens and strobe lights activated at irregular intervals are more effective than either a constant visual display or loud noises generated at fixed intervals. Supplementing these techniques with occasional lethal measures (e.g., shooting) further enhances their continued efficacy. Properly trained dogs confined within the boundaries of the protected resource are very effective at frightening wildlife.

## Visual Devices

### Reflectors

**Function:** Reflectors are used to frighten wildlife away from resources by reflecting flashes of natural or artificial light.

**Description:** Reflectors are passive devices that can be used alone or in conjunction with artificial lights to frighten wildlife species away from protected areas. Mylar reflecting tape can be attached to stakes or suspended from wire strung between stakes placed within and around the protected area. The mylar tape is designed to reflect sunlight. Movement by the wind causes sporadic flashes and noise. Wrinkled aluminum foil also works well as an effective reflector, producing sporadic flashes of light and erratic noises when blown by the wind.

Reflectors fitted within molded mounts attached to stakes are more permanent and come ready to install. These types of reflectors are most beneficial to increase the area of coverage or efficacy of artificial light systems operated at night.

**Operation:** Mounted reflectors can be inserted directly into the ground. Otherwise, install posts or hang reflectors from vegetation or debris in the area. Concentrate reflectors on areas where the target species is most active. Reflectors should not be set in shaded areas. When reflectors are set to complement artificial light, ensure that a clear path exists between the light source and the reflector. Occasionally alter reflector locations to reduce habituation.

**Advantages:** Reflectors are relatively inexpensive and easy to install, requiring a minimal amount of maintenance. They are not hazardous to wildlife or humans.

**Disadvantages:** Ineffective at night or on cloudy days, unless combined with artificial light. Most animals habituate to reflectors.

### Sources:

C. Frensch, Ltd.  
Orchard Supply Company  
Strieter Corporation  
Ted Dodge Service  
Trucall

## Artificial Light

**Function:** Light can be used to illuminate areas or to produce flashes to frighten animals away from protected areas. Nocturnal species avoid well-lit areas, while other species may restrict their use of these sites. Light flashes are used to scare animals away from specific areas.

**Description:** Light sources vary from simple to complex. All, however, require a lamp, power source, and protective cover. Several lamps can be operated by a common power source. Long-term efficacy of sporadic lights can be enhanced by adding several optional devices, such as strobes, timers, revolving platforms, or trip-switches.

**Operation:** Continuous lights mounted on poles need to be strategically placed so the entire area is well lit. Sporadic lights need to be set so they are clearly visible by intruding animals, preferably so the flash will illuminate the animal. Changing locations and schedule of periodic flashes helps to reduce habituation. Artificial lights need to be checked frequently to ensure that the light and optional mechanisms are functioning properly.

**Advantages:** Lights are active systems and do not depend on natural elements (e.g., sunlight, wind) to operate. Lights are not hazardous to wildlife or humans. Nocturnal species do not habituate to continuous light. Further, animals are less likely to habituate to actively changing systems than to passive, constant systems.

**Disadvantages:** Lights can be expensive and require more effort to install and maintain than passive systems. Lights can be distracting and

bothersome to neighbors. As animals habituate to light systems, lights become less effective.

**Sources:**

Bird-X, Inc.  
The Huge Company, Inc.  
Tripp-Lite Manufacturing Company

## Effigies

**Function:** Human or predator effigies are installed to create the illusion of an unsafe environment.

**Description:** Effigies create the illusion that natural predators or humans are active in the protected area. Predominately used to deter birds, effigies are also installed to frighten big game. Scarecrows (e.g., human effigies) can be anything from shirts stuffed with straw and fastened to poles to lifelike mechanized models. Bright colors, or reflectors and streamers that flutter in the wind, increase the efficacy. Predator effigies also encompass the continuum from simple silhouettes to lifelike models. The more realistic models, and effigies that incorporate some movement, are the most effective. Human effigies paired with occasional shooting can be very effective.

**Operation:** Select effigies that are likely to produce an alarming or threatening situation. For example, raptor models may induce avoidance by small birds but go unheeded by elk. Place effigies where they are clearly visible to the target species. Change locations periodically to reduce habituation.

**Advantages:** Simple models are inexpensive and easy to install. Effigies are not dangerous to wildlife or humans.

**Disadvantages:** Complex models can be relatively expensive and may require some maintenance. Efficacy of effigies declines as animals habituate to them

unless the devices are paired with other negative consequences.

**Sources:**

Bird-X, Inc.  
Flambeau Products Corporation  
FLR, Inc.  
The Huge Company, Inc.  
Orchard Supply Company  
Sutton Ag Enterprises, Inc.

## Balloons, Kites, and Flags

**Function:** Balloons, kites, and flags are used to scare wildlife away from an area by creating an unfamiliar situation.

**Description:** Balloons (Figure 35) come in a variety of colors and sizes. Several models are painted with bright colors and large circular patterns to resemble the eyes and open mouth of predators. Balloons are either suspended from poles or filled with helium and tethered to stakes.

Kites are hung from wires or strings strung across the site, or flown over the site by an operator. Kites designed to frighten birds may double as predator effigies. A predator design, combined with the fluttering and dipping motions of a kite, can be realistic and effective.

The flapping motions and noises created by flags can also restrict animal use of an area. Flags can be purchased, or constructed by attaching sheets of plastic or cloth to a stake. Fabrics with distinct and clearly visible colors should be used to create flags.

**Operation:** Balloons, kites, and flags have to be clearly visible to be effective. The recommended number of balloons and flags depends on the situation, though as few as one per acre may be adequate. Switching colors and locations of balloons and flags can prolong the effective life of these devices. A skilled operator can effectively cover a substantial area with a single kite.




Figure 35—"Scare-Eye" balloon.

**Advantages:** Initial costs of balloons, kites, and flags are relatively inexpensive. These devices are easy to install and require minimum amounts of maintenance. None of these devices are detrimental to wildlife or the environment.

**Disadvantages:** Operator-flown kites can become labor intensive. Helium-filled balloons need to be maintained on a regular basis. The efficacy of these devices declines as animals habituate to them over time.

**Sources:**

Atmospheric Instrumentation  
Research, Inc.  
Birdbusters  
Bird Scare Predator Eye, Inc.  
Bird-X, Inc.  
C. Frensch, Ltd.  
Cochranes of Oxford, Ltd.  
Ecopic  
Hartman's Plantation, Inc.  
Margo Supplies, Ltd.  
Nixalite of America, Inc.  
Orchard Supply Company  
Reed-Joseph International Company  
Rid-A-Bird, Inc.  
Sutton Ag Enterprises, Inc.  
Wildlife Control Technology, Inc.

## Noisemakers

### Distress Calls, Sirens, and Horns

**Function:** Distress calls mimic the sounds of birds in trouble, which frightens and discourages birds from landing in an area. Sirens, horns, and other sounds can also be used to scare birds and mammals from an area.

**Description:** Distress calls are commonly generated by battery-operated portable cassette recorders. Recorders are connected to an amplifier and a long-range speaker. Recordings of distress calls from a variety of birds are commercially available. Predator calls, barking dogs, or human noises can also be recorded and played over the same system to scare problem species.

A variety of compressed air or battery-operated sirens and horns are commercially available (Figure 36). Like the cassette recorders, these are attached to a directional speaker that focuses the noise toward the problem area.

**Operation:** Place the cassette player with the directional speaker pointed toward the major problem area. Turn the volume to a level loud enough for birds to hear the distress calls over other noises. Initiate calls at times when the birds normally gather to feed; continue the distress calls until depredation ceases. Sounds to scare other animals can be generated in a similar fashion. Habituation to the sounds can be delayed if the machine is not operated continuously and tapes are switched periodically. A trip device or timer set to operate the player sporadically will further reduce the rate of habituation.

Sirens and horns are operated similarly to distress calls. Strategically place the siren or horn where it will be directed at the most severe depredation sites. A timer or trip device attached to the siren

or horn will enhance the efficacy of the operation. Hand-held sirens and horns can also be used by an operator to scare animals away from a site.

**Advantages:** Portable cassette players can easily be installed or moved to new locations. After a moderate initial expenditure, the operating expenses are minimal. Distress calls or other vocalizations are not harmful to wildlife.

**Disadvantages:** Animals generally habituate to sounds that are not reinforced with punishment. Hand-operated devices can become labor intensive. Loud noises are often distracting to neighbors. Electronic or battery-operated devices placed in the field require weatherproofing and frequent maintenance.

#### Sources:

Birdbusters  
Bird-X, Inc.  
C. Frensch, Ltd.  
Denver Wildlife Research Center  
Falcon Safety Products, Inc.  
Farm and Industrial Supplies Company  
Johnny Stewart  
Margo Supplies, Ltd.  
P.S. Olt Company  
Reed-Joseph International Company

Signal Broadcasting Products, Inc.  
Tomko Enterprises, Inc.  
Weitech, Inc.

### Ultrasonic Devices

**Function:** Ultrasonic devices (Figure 37) emit high frequency sounds to discourage wildlife from entering an area.

**Description:** Ultrasonic frequencies exceed 20,000 hertz, a level that is not audible or disturbing to humans. Several models of ultrasonic devices are commercially available. Devices generally consist of an ultrasonic unit and a battery or solar power source. Some devices are equipped with mounting brackets; others are affixed to poles. Optional motion detectors are available that activate the device for a specified time interval whenever it is approached by an animal.

**Operation:** Install the ultrasonic devices such that they cover or at least encircle the problem area. Spacing will depend on the particular device; most are reported effective up to at least 30 feet. Timers or trip devices will prolong the power supply.




Figure 36—"Bird Gard ABC" noisemaker system used to frighten animals.




Figure 37—"Usonic Sentry" ultrasonic device used to frighten animals.

**Advantages:** Ultrasonic devices are relatively inexpensive to operate. Animals do not habituate to ultrasonic frequencies as readily as they do to audible sounds. Humans are not distracted by ultrasonic frequencies and the devices are not detrimental to the environment or harmful to animals.

**Disadvantages:** The original expenditure for ultrasonic devices to cover a large area can be expensive. Ultrasonic frequencies diminish rapidly with increasing distance from the source. Further, shadows of ultrasonic frequencies are formed wherever sound waves are obstructed. Areas with debris or substantial amounts of vegetation will need to be cleared or additional devices installed. Even though ultrasonic frequencies are inaudible, high decibel levels may damage human hearing; appropriate hearing protection should be provided to operators.

**Sources:**

Adams Dominion, Inc.  
Bird-X, Inc.  
C. Frensch, Ltd.  
Falcon Safety Products, Inc.  
Farm and Industrial Supply Company

Johnny Stewart  
Margo Supplies, Ltd.  
Medline of Colorado  
Orchard Equipment and Supply  
Reed Joseph International Company  
Signal Broadcast Products, Inc.  
Weitech, Inc.

## Automatic Exploders

**Function:** Automatic gas exploders (Figure 38) emit loud explosive noises to frighten animals away from an area.

**Description:** Automatic gas exploders operate on propane or acetylene. The exploder operates by igniting gas collected within a chamber. A cannon or tube directs and amplifies the explosive noise. An adjoining storage tank supplies gas to the exploder. Most tanks contain adequate fuel for 12,000 or more explosions. The exploder is often mounted on a tripod to elevate it above surrounding vegetation. Exploders come with various options including timers, rotating directional mechanisms, automatic erratic volume control, dual and triple cannons, and attachable visual displays.

**Operation:** Spread exploders across the problem area. The necessary number of exploders depends on your objectives and site-specific factors; however, operational estimates for the coverage of one exploder range from 3 to 40 acres. Field observations will indicate how many and where the exploders need to be established. Elevate or set exploders so that the

sound will not be muffled by surrounding vegetation or debris. Habituation will be less rapid if exploders are fired in irregular directions, and with varying time intervals and magnitudes. For best results, move the exploders to new locations every other day.

**Advantages:** Automatic cannons work in the absence of an operator and require only moderate maintenance. The exploders are not harmful to wildlife.

**Disadvantages:** Animals habituate to the noise after varying periods of time. Labor to move exploders repeatedly can become expensive if a large number of exploders are necessary to cover an extensive area. The noise may be a problem for neighbors.




Figure 38—Assorted "Zon Scarecrow Guns" used to frighten animals.



**Sources:**

Agricultural Supply, Inc.  
C. Frensch, Ltd.  
Field Systems Corporation  
Harmon International, Inc.  
H.C. Shaw Company  
Margo Supplies, Ltd.  
M.J. Flynn, Inc.  
Orchard Equipment and Supply Co.  
Pisces Industries  
Reed-Joseph International Company  
Sutton Ag Enterprises, Inc.  
Teiso Kasei Company  
Wildlife Control Technology, Inc.

## Pyrotechnic Devices

**Function:** Hand-fired noise devices projected by an operator over an area to frighten animals.

**Description:** Pyrotechnic devices (Figure 39) are fired from a hand-held gun by an operator. Various types are available, but most fall within the general categories of bangers, screamers and shell crackers. Bangers can be projected from 75 to 125 feet and explode with a loud bang or pop. Screamers make a loud screeching or whistling noise as they travel 300 to 350 feet across a protected site. At night, screamers produce a visual effect similar to a twirling bottle rocket. Banger and screamer cartridges are projected by .22-caliber blanks fired from hand-held revolvers. Shell crackers, the loudest of the pyrotechnic devices, explode with a bang. Fired from an open choke 12-gauge shotgun, the shell cracker will travel about 200 feet.

**Operation:** An operator makes regular visits to a depredation site, firing pyrotechnics across the site whenever the offending species is present. Devices should be projected such that they

explode over the site and not so that they hit animals or explode on the ground. Direct hazing should follow pyrotechnics whenever animals fail to respond to the noise. Possession and use of pyrotechnics may require a permit from the fire marshal. Check the launch barrel after each discharge because wads may stick in the barrel.

**Advantages:** Animals are less likely to habituate to pyrotechnics because they are delivered at specific times and directed toward specific places by an operator. Animals that fail to heed pyrotechnic devices can be immediately hazed. Noise produced by pyrotechnic devices is less offensive to neighbors because devices are fired only when necessary. Devices are relatively safe for both wildlife and humans.

**Disadvantages:** Pyrotechnics require the presence of an experienced operator to monitor sites and to fire them. Pyrotechnics can cause fires and are hazardous to the operator if not handled with care.

**Sources:**

Agricultural Supply, Inc.  
C. Frensch, Ltd.  
Margo Supplies, Ltd.  
Marshall Hyde, Inc.  
New Jersey Fireworks Company  
Reed-Joseph International Company  
Stoneco, Inc.  
Sutton Ag Enterprises, Inc.  
Wald & Company  
Western Fireworks Company  
Wildlife Control Technology, Inc.



Figure 39—Pyrotechnic devices used to frighten animals.

## Alternative Forage

**W**ildlife damage to forest resources is invariably induced by foraging animals. Wildlife feed on seedlings and the sapwood of larger trees to obtain a source of nourishment. Dietary selection, however, is relative and depends on the available options. Providing wildlife with viable alternative foraging options can alleviate the extent of foraging pressure directed toward forest resources.

Several approaches can be used to provide wildlife with alternative foods. Desirable foods can be distributed across problem areas or on adjacent sites to encourage animals away from the protected resources. For example, alfalfa distributed along migratory trails may reduce the tendency of ungulates to browse seedlings. Another approach is to plant or encourage the establishment of natural forages that are preferred by wildlife. Some plants, such as cat's ear, are ingested by most herbivorous rodents, as well as ungulates. Animals can also be provided food supplements in semipermanent structures strategically placed adjacent to or within sites that are being damaged or that are vulnerable to damage.

Before implementing a feeding program to reduce forest damage, the long-term consequences need to be considered. Alternative forages can increase or prolong the presence of wildlife on selected sites. Increased resources may encourage additional animals to frequent the area, or an improved nutritional status may enhance reproductive success. Further, resource-dependent territorial boundaries of rodents may shrink with improved forage resources, which in turn permits more individuals to exist within a given forest unit. Big-game herds may suspend or delay migratory movements to utilize available forages. A feeding program, therefore, might actually increase wildlife pressure on forest resources if the program is not sustained, or fails to meet the increasing demands.

A successful program needs to be specific in targeting a problem for a known time interval. In addition, a means to continue the program indefinitely or a means to wean the supplemented animal from the program should be identified. The potential for animals to revert to protected forest resources also needs to be anticipated and avoided.

The supplemental bear feeding program coordinated by the Washington Forest Protection Association is an example of an effective feeding program to prevent forest damage. The program provides a favorable alternative to bears. They are offered a choice of ingesting nutritious pellets rather than girdling trees when other natural foods are limited. A natural weaning from the program occurs as bears revert to natural forages (e.g., berries) as they become available.

### Supplemental Bear Feeding Program

**Function:** Bears are provided a supplemental food to reduce their destructive foraging on trees during the spring.

**Description:** Self-feeders are constructed from 55-gallon drums (Figure 40). Bears can obtain pellets from an opening cut in the lower front portion of the barrel. The feeding plate is separated from the storage compartment by a metal sheet inserted diagonally within the barrel. Pellets taken from the feeding plate are automatically replaced as pellets

fall through a narrow gap beneath the metal sheet. This self-feeding mechanism permits a continuous supply of pellets, but prohibits bears from spilling or playing with the food. A heavy roof insulated with foam keeps the pellets dry and limits bears to feeding from the front entrance.

The supplemental pellets are produced under the direction of the Washington Forest Protection Association. Sugars are considered the most important ingredients to alleviate damage to trees. Pellets also contain fats, proteins, vitamins, and minerals to provide a balanced diet for bears. A low moisture content (less than 10%) enhances bear acceptance and increases the viable shelf life of pellets. Individual pellets resemble a greenish-colored dry dog food and are 0.25 inches in diameter and about 0.5 inches long.

**Operation:** Supplemental feed is provided only in the spring and early summer when bears are foraging on trees. Feeding stations are installed in sites as close as possible to current or anticipated damage. Preferred sites are close to roads to facilitate stocking feeders with pellets, but hidden from public view to avoid poaching. Sites are kept away from high public use areas to avoid bear-human conflicts.




Figure 40—Bear feeding station used to supply bears with supplemental pellets during the spring.

Bear feeders are securely fastened to trees approximately 10 inches off the ground to avoid water and rodent problems. Bait (e.g., a beaver carcass) may be used to initially attract bears to stations, but is not necessary once bears begin feeding. Each station holds approximately 200 pounds of pellets. Active stations need to be restocked once a week. Wet or powdery feed is

removed before a station is restocked with new pellets. At the end of the damage season all feeding stations are removed from the forests.

**Advantages:** Supplemental feeding of bears is not detrimental to bears or other wildlife species and is usually well received by the general public.

**Disadvantages:** The program is expensive and labor intensive to implement and maintain. The long-term impacts of supplemental feeding on subsequent bear behavior, reproductive success, and tree damage are not well understood.

**Source:**  
Washington Forest Protection Assoc.

---

## Sources

Adams Dominion, Inc.  
P.O. Box 690  
Prospect, KY 40059  
Phone: (502) 228-3361

ADPI Enterprises, Inc.  
3621 B Street  
Philadelphia, PA 19134  
Phone: 800-621-0275  
Phone: (215) 425-8866  
Fax: (215) 739-8480

Agricultural Supply, Inc.  
1435 Simpson Way  
Escondido, CA 92029  
Phone: 800-527-6699  
Fax: (619) 741-9412

American Feed & Farm  
3310 H Street  
Omaha, NE 68107  
Phone: 800-642-9904

American Forestry Technology, Inc.  
1001 North 500 West  
West Lafayette, IN 47906  
Phone: (317) 583-3311  
Fax: (317) 583-3318

Animal Care Equipment & Services  
P.O. Box 3275, 613 Lee Bert Way  
Crestline, CA 92325  
Phone: 800-338-2237  
Phone: (909) 338-1791  
Fax: (909) 338-2799

Animal Management, Inc.  
720 Eppley Road  
Mechanicsburg, PA 17055-9786  
Phone: 800-745-8173  
Phone: (717) 790-9347  
Fax: 800-745-8193

Animal Repellents, Inc.  
P.O. Box 510  
Orchard Hill, GA 30266  
Phone: 800-241-5064  
Phone: (404) 227-8222

APGAR, INC., Inc.  
Mill River Supply  
375 Adams  
Bedford Hills, NY 10507  
Phone: (914) 666-5774  
Fax: (914) 666-9183

Archem Corporation  
1514 11th Street  
Portsmouth, OH 45662  
Phone: 800-635-1125  
Fax (614) 353-1124

Atmospheric Instrumentation  
Research (Air), Inc.  
8401 Baseline Road  
Boulder, CO 80303  
Phone: (303) 443-7187  
Fax: (303) 499-1767

Avitrol Corporation  
7644 East 46th Street  
Tulsa, OK 74145  
Phone: 800-633-5069  
Phone: (918) 622-7763  
Fax: (918) 622-2527

Baygard-Bay Mills, Ltd.  
6 Holtby Avenue  
Brampton, Ontario  
Canada L6X 2M1  
Phone: (905) 456-3394  
Fax: (905) 456-1452

Bell Laboratories, Inc.  
3699 Kinsman Boulevard  
Madison, WI 53704  
Phone: (608) 241-0202  
Fax: (608) 241-9631

Bernardo Chemical, Ltd., Inc.  
P.O. Box 9040  
Rock Mount, NC 27804-7040  
Phone: (919) 443-9325  
Fax: (919) 443-5386

B&G Chemicals & Equipment Co., Inc.  
10539 Maybank  
Dallas, TX 75354-0428  
Phone: 800-345-9387  
Phone: (214) 357-5741  
Fax: (214) 357-4514

Bigelow Trap Company  
979 Milford Avenue  
Marysville, OH 43040  
Phone: (513) 642-6786

Birdbusters  
1083 Thomas Jefferson Street NW  
Washington, DC 20007  
Phone: 800-662-4737  
Phone: (202) 338-6263  
Fax: (202) 338-6268

Bird Scare Predator Eye, Inc.  
132 East Demont Avenue, #122  
Little Canada, MN 55117-1561  
Phone: 800-252-0933  
Phone: (612) 482-1123  
Fax: (612) 482-1241

Bird-X, Inc.  
730 West Lake Street  
Chicago, IL 60661  
Phone: 800-662-5021  
Phone: (312) 648-2191  
Fax: (312) 648-0319

B-Kind Animal Control Equipment  
Southeastern Metal Products, Inc.  
1200 Foster Sreet NW, P.O. Box 93038  
Atlanta, GA 30377  
Phone: (404) 351-6686

Blue Mountain Industries  
20 Blue Mountain Road, P.O. Box 4000  
Blue Mountain, AL 36204  
Phone: (205) 237-9461  
Fax: (205) 237-8816

B.N. Bolton, Inc.  
Route 1, Site 11, Comp. 42  
Vernon, British Colombia  
Canada V1T 6L4

Boehringer Ingelheim Animal Health, Inc.  
Anchor Division  
2621 North Belt Highway  
St. Joseph, MO 64506-2002  
Phone: (816) 233-1385

Bonide Products, Inc.  
2 Wurz Avenue  
Yorkville, NY 13495  
Phone: (315) 736-3231  
Fax: (315) 736-7582

Brookside Industries, Inc.  
Route 1, P.O. Box 158  
Tunbridge, VT 05077  
Phone: (802) 889-3737

Burlington Scientific Corporation  
222 Sherwood Avenue  
Farmingdale, NY 11735  
Phone: (516) 694-9000  
Fax: (516) 694-9177

Bushwacker Backpack & Supply Co., Inc.  
P.O. Box 4721  
Missoula, MT 59806  
Phone: 800-695-3394  
Phone: (406) 728-6241  
Fax: (406) 728-8800

C. Fensch, Ltd.  
168 Main Street, P.O. Box 67  
Grimsby, Ontario  
Canada L3M 4G1  
Phone: (905) 945-3827  
Fax: (905) 945-4128

Champon 100% Natural Products, Inc.  
10528 Mendocino Lane  
Boca Raton, FL 33428  
Phone: (407) 852-4143  
Fax: (407) 852-4143

The Chas. H. Lilly Company  
P.O. Box 83179  
Portland, OR 97283-0179  
Phone: (503) 289-5937  
Fax: (503) 289-9216

C.J. Martin Company  
P.O. Box 630009, 600 West Main Street  
Nacogdoches, TX 75963  
Phone: (409) 560-8282  
Fax: (409) 560-8331

Cochranes of Oxford, Ltd.  
Leaffield Witney  
England, UK OX8 5NY  
Phone: 099387-641

Conwed Plastics  
2810 Weeks Avenue SE  
Minneapolis, MN 55414  
Phone: 800-426-6933  
Phone: (612) 623-1700  
Fax: (612) 623-2500

"Coon Getter" Cage Traps & Animal Lures  
Rural Route 4, P.O. Box 109  
Miller, SD 57362  
Phone: (605) 853-2545

Coughlan Products Corporation  
P.O. Box 1888  
Paterson, NJ 07509  
Phone: (201) 742-6468  
Fax: (201) 742-0359

Critter Control Wildlife Mgmt. Supplies  
640 Starkweather  
Plymouth, MI 48170  
Phone: 800-451-6544  
Phone: (313) 453-6300

Critter Gitters  
4317 Nancy Creek Road  
P.O. Box 462  
Kettle Falls, WA 99141  
Phone: (509) 738-2230

Dairyland Power Fence Company  
North 3985 Hidden Valley Road  
Hatley, WI 54440  
Phone: (715) 446-2297

Davenport Seed Corporation  
P.O. Box 187  
Davenport, WA 99122  
Phone: 800-828-8873  
Phone: (509) 725-7015

Daviyn Manufacturing Company, Inc.  
P.O. Box 49  
Spring City, PA 19475  
Phone: (215) 948-5050  
Fax: (215) 948-5454

Degesh America, Inc.  
P.O. Box 116  
275 Triangle Drive  
Weyers Cave, VA 24486  
Phone: (703) 234-9281  
Fax: (703) 234-8225

Denver Wildlife Research Center  
USDA-APHIS, Animal Damage Control  
P.O. Box 25266, Building 16  
Denver Federal Center  
Denver, CO 80225-0266  
Phone: (303) 236-7820

Design Packaging  
2851 North Sunnyside, #106  
Fresno, CA 93727  
Phone: (209) 292-9363  
Fax: (209) 298-9194

Don Sprague Sales, Inc.  
1470 Aztec  
Woodburn, OR 97071  
Phone: 800-423-0158  
Phone: (506) 981-0297

Douglas Products & Packaging Company  
1500 East Old 210 Highway  
Liberty, MO 64068  
Phone: 800-223-3684  
Fax: (816) 781-1043

Dragon Corporation  
P.O. Box 7311  
Roanoke, VA 24019  
Phone: 800-533-2488  
Phone: (703) 362-3657  
Fax: (703) 362-9171

Dr. T's Nature Products Company, Inc.  
P.O. Box 682, U.S. 19 North  
Pelham, GA 31779  
Phone: 800-299-6288  
Phone: (912) 294-9742

Duke Company  
508 Brame Avenue, P.O. Box 555  
West Point, MS 39773  
Phone: (601) 494-6767  
Fax: (601) 494-5360

Earl May Seed & Nursery Company  
208 North Elm Street  
P.O. Box 500  
Shenandoah, IA 51603  
Phone: (712) 246-1020

Eckroat Seed Company  
1106 M.L. King Avenue  
P.O. Box 17610  
Oklahoma City, OK 73136  
Phone: 800-331-7333  
Phone: (405) 427-2489

Eco-Bait  
Donald D. Hicks  
P.O. Box 321  
Geraldine, MT 59446  
Phone: (406) 737-4291

Ecopic  
725 S. Adams Road, Suite 270  
Birmingham, MI 48009  
Phone: (810) 647-0505  
Fax: (810) 647-7811

E.I. Dupont DeNemours & Company, Inc.  
Polymer Products Department  
Wilmington, DE 19898  
Phone: (302) 774-1000  
Fax: (302) 774-7321

Faesy & Besthoff, Inc.  
143 River Road, P.O. Box 29  
Edgewater, NJ 0702  
Phone: (201) 945-6200  
Fax: (201) 945-6145

Falcon Safety Products, Inc.  
25 Chubb Way, P.O. Box 1299  
Somerville, NJ 08876-1299  
Phone: (908) 707-4900  
Fax: (908) 707-8855

Farm and Industrial Supply Company  
P.O. Box 31510  
Stockton, CA 95213  
Phone: 800-221-2884

Farnam Company, Inc.  
Sudbury Consumer Products Division  
P.O. Box 34820  
Phoenix, AZ 85067-4820  
Phone: 800-825-2555  
Phone: (602) 285-1660  
Fax: (602) 285-1803

F.B.N. Plastics  
225 North L Street  
Tulare, CA 93274  
Phone: (209) 688-7269

Ferrell and Company  
P.O. Box 92  
Union, MS 39365  
Phone: 800-848-0759  
Phone: (601) 774-8983

Ferrett Laboratories, Inc.  
P.O. Box 437  
Sliger, WA 53086  
Phone: (414) 644-5252

Field Systems Corporation  
P.O. Box 1374  
Huron, SD 57350  
Phone: (605) 352-7703

Fiambeau Products Corporation  
15981 Balplast Road  
Middlefield, OH 44062  
Phone: (216) 632-1631  
Fax: (216) 632-1581

FLR, Inc.  
P.O. Box 108  
Midnight, MS 39115  
Phone: (601) 247-4409

Forest Protection Products  
P.O. Box 1047  
Coos Bay, OR 97420  
Phone: 800-289-7659  
Phone: (503) 267-2622  
Fax: (503) 269-7300

Forestry Suppliers, Inc.  
205 West Rankin Street, P.O. Box 8397  
Jackson, MS 39284-8397  
Phone: 800-647-5368  
Fax: (601) 355-5126

Fort Dodge Chemical Company  
P.O. Box 2021  
Lompoc, CA 93438  
Phone: (805) 736-0065

Fuhrman Diversified, Inc.  
2912 Bayport Boulevard  
Seabrook, TX 77586-1501  
Phone: (713) 474-1388  
Fax: (713) 474-1390

Gallagher Power Fence, Inc.  
P.O. Box 708900  
San Antonio, TX 78270-8900  
Phone: 800-553-1680  
Phone: (512) 494-5211  
Fax: (507) 864-7054

Geotek, Inc.  
P.O. Box 607, 13433 Highway 52 SW  
Chatfield, MN 55923  
Phone: (507) 867-4243  
Fax: (507) 867-3713

Geyer Irrigation  
700 5th Street  
Arbuckle, CA 95912  
Phone: (916) 476-2253  
Fax: (916) 476-2256

GMK Company  
43 Pineview Avenue, B18  
Cardiff, NJ 08232  
Phone: (609) 641-8490

Goodart's  
P.O. Box 1466  
Hemphill, TX 75948  
Phone: 800-458-6608

Grassland Supply  
Route 3, P.O. Box 6  
Council Grove, KS 66846  
Phone: 800-527-5487  
Phone: (316) 767-5487  
Fax: (316) 767-6679

Grawes Animal Lures  
P.O. Box 306  
Wahpeton, ND 58074-0306  
Phone: (218) 643-3292

Green Valley Blueberry Farm  
9345 Ross Station Road  
Sebastopol, CA 95472  
Phone: 800-827-9590  
Phone: (707) 887-7496  
Fax: (707) 887-7499

Gremar Company  
4719 Cody Drive  
P.O. Box 65003  
West Des Moines, IA 50265  
Phone: (515) 263-3007  
Fax: (515) 221-9243

Guardian Trap Company  
P.O. Box 1935  
San Leandro, CA 94577  
Phone: (415) 357-0900

Gustafson, Inc.  
P.O. Box 660065  
Dallas, TX 75266-0065  
Phone: 800-527-4781  
Phone: (214) 985-8877  
Fax: (214) 985-1696

HACO, Inc.  
537 Atlas Avenue, P.O. Box 7190  
Madison, WI 53707  
Phone: (608) 221-6200  
Fax: (608) 221-6208

Hadley Products Company  
Route 1  
Marietta, OH 45750  
Phone: (614) 373-9298

Hancock Trap Company  
P.O. Box 268  
Custer, SD 57730  
Phone: (605) 673-4128

Harmon International, Inc.  
P.O. Box 1827  
Minot, ND 58702  
Phone: (701) 839-6717  
Fax: (701) 839-8679

Hartman's Plantation, Inc.  
310 60th Street, P.O. Box E  
Grand Junction, MI 49056-0805  
Phone: (616) 253-4281  
Fax: (616) 253-4457

H.B. Sherman Traps, Inc.  
P.O. Box 20267  
Tallahassee, FL 32316  
Phone (904) 562-5566

H.C. Shaw Company  
4554 Quantas Lane, Suite 1  
Stockton, CA 95206  
Phone: 800-221-2884  
Phone: (209) 983-8484  
Fax: (209) 983-8449

Herd Seeder Company  
P.O. Box 448  
Logansport, IN 46947  
Phone: (209) 753-6311  
Fax: (209) 722-4106

Hoffman Trapping Supplies, Ltd.  
P.O. Box 805  
Russell, Manitoba  
Canada ROJ 1W0  
Phone: (204) 773-2270

The Huge Company, Inc.  
7625 Page Boulevard  
St. Louis, MO 63130  
Phone: 800-873-4843  
Phone: (314) 725-2555

IntAgra, Inc.  
8500 Pillsbury Avenue South  
Minneapolis, MN 55420-2219  
Phone: 800-468-2472  
Phone: (612) 881-5535  
Fax: (612) 881-7002

Integral Designs  
3628 Burnsland Road SE  
Calgary, Alberta  
Canada T2M 4L5  
Phone: (403) 285-9731

Internet, Inc.  
2730 Nevada Avenue North  
Minneapolis, MN 55427  
Phone: 800-328-8456  
Phone: (612) 541-9690  
Fax: (612) 541-9692

J.A. Cissel Manufacturing Company  
P.O. Box 2025  
Lakewood, NJ 08701  
Phone: 800-631-2234  
Fax: (908) 901-1166

J.C. Ehrlich Chemical Company  
500 Spring Ridge Drive  
Reading, PA 19612  
Phone: 800-488-9495  
Phone: (215) 372-9700  
Fax: (215) 378-9744

J/J Fur Shop  
717 North 23rd Street  
Hot Springs, SD 57747  
Phone: (605) 745-5114

Johnny Stewart  
5100 Fort Avenue, P.O. Box 7594  
Waco, TX 76714-7594  
Phone: (817) 772-3261  
Fax: (817) 772-3670

J.T. Eaton & Company, Inc.  
1393 East Highland Road  
Twinsburg, OH 44087  
Phone: 800-321-3421  
Phone: (216) 425-7801  
Fax: (216) 425-8353

Kemi-K Products, Inc.  
P.O. Box 21  
Yakima, WA 98907  
Phone: 800-321-5354  
Phone: (509) 453-1314

Ketch-All Company  
4149 Santa Fe Road, #2  
San Luis Obispo, CA 93401  
Phone: (805) 543-7223  
Fax: (805) 543-7145

Keystone Steel & Wire  
7000 SW Adams Street  
Peoria, IL 61641  
Phone: 800-447-6444  
Phone: (309) 697-7422

K Fence Systems  
Route 1, P.O. Box 195  
Zumbro Falls, MN 55991  
Phone: (507) 753-2943  
Fax: (507) 753-2706

Kiwi Fence Systems  
1145 East Roy Furman Highway  
Waynesburg, PA 15370-8070  
Phone: (412) 627-8158  
Fax: (412) 627-9791

Kness Manufacturing Company, Inc.  
Highway 5 South, P.O. Box 70  
Albia, IA 52351-0070  
Phone: 800-247-5062  
Phone: (515) 932-7846  
Fax: (515) 932-2456

Krofick Outdoor Supply  
30 Lightcap Road  
Latrobe, PA 15650  
Phone: (412) 537-7923

Laird Plastics, Inc.  
6311 Erdman Avenue  
Baltimore, MD 21205-3585  
Phone: (410) 485-9100  
Fax: (410) 485-3033

Lakeshore Enterprises  
2804 Benzie Highway  
Benzonia, MI 49616  
Phone: (616) 882-9601

Leggett's Supplies  
7308 Monroe Road  
Boonsboro, MD 21713  
Phone: (310) 432-6210  
Fax: (301) 432-8480

LiphaTech, Inc.  
3600 West Elm Street  
Milwaukee, WI 53209  
Phone: 800-558-1003  
Phone: (414) 351-1476  
Fax: (414) 351-1847

Live Wire Products, Inc.  
1127 E Street  
Marysville, CA 95901  
Phone: 800-272-9045  
Phone: (916) 743-9045  
Fax: (916) 743-0609

Manufacturing Systems, Inc.  
Tru-Catch Traps  
300 Industrial Street, P.O. Box 816  
Belle Fourche, SD 57717-0816  
Phone: 800-247-6132  
Phone: (605) 892-2717  
Fax: (605) 892-6327

Margo Supplies, Ltd.  
P.O. Box 5400  
High River, Alberta  
Canada T1V 1M5  
Phone: (403) 652-1932  
Fax: (403) 652-3511

Marshall Hyde, Inc.  
P.O. Box 497  
Port Huron, MI 48060  
Phone: (313) 982-2140

McGill Metal Products Company  
142 East Prairie Street, P.O. Box 177  
Marengo, IL 60152  
Phone: (815) 568-7244

Medlinc of Colorado  
623 Broken Spoke Road  
Grand Junction, CO 81054  
Phone: (303) 241-0366  
Fax: (303) 241-0366

Micro-Gen Equipment Corporation  
10700 Sentinel Drive  
San Antonio, TX 78217  
Phone: 800-777-8570  
Phone: (210) 654-8570  
Fax: (210) 654-3613

Miller Chemical and Fertilizer Corporation  
P.O. Box 333, Radio Road  
Hanover, PA 17331  
Phone: 800-233-2040  
Phone: (717) 632-8921  
Fax: (717) 632-4581

Miller Net and Twine  
P.O. Box 18787, 1674 Getwell Road  
Memphis, TN 38181-0787  
Phone: 800-423-6603  
Fax: (901) 743-6580

Minnesota Plasti-Catch  
P.O. Box 43  
Darwin, MN 55234  
Phone: (612) 275-2523

Minnesota Trapiine Products  
6699 156th Avenue NW  
Pennock, MN 56279  
Phone: (612) 599-4176  
Fax: (612) 599-4314

Mississippi Valley Forest Products, Inc.  
P.O. Box 1128  
Dubuque, IA 52004  
Phone: (319) 583-2795

Mitlying Development  
P.O. Box 43  
Darwin, MI 55324  
Phone: (612) 275-2523

M.J. Flynn, Inc.  
6408 Collamer Road  
East Syracuse, NY 13057-1032  
Phone: (315) 437-6536  
Fax: (315) 432-1315

M&M Fur Company  
P.O. Box 15  
Bridgewater, SD 57319-0015  
Phone: 800-658-5554  
Phone: (605) 729-2535  
Fax: (605) 729-2536

Monterey Chemical Company  
5150 North 6th Street  
P.O. Box 5317  
Fresno, CA 03755  
Phone: (209) 225-4770  
Fax: (209) 225-1310

Montgomery Traps, Inc.  
P.O. Box 121  
Mahaffey, PA 15757  
Phone: (814) 277-6653

Morrison Manufacturing Corporation  
Highway 175, P.O. Box 52  
Morrison, IA 50657  
Phone: 800-648-CAGE  
Phone: (319) 345-6406  
Fax: (319) 345-6429

Motomco, Ltd.  
3699 Kinsman Boulevard  
Madison, WI 53704  
Phone: (813) 447-3417  
Fax: (608) 241-2238

Naltex Plastics, Inc.  
220 East St. Elmo, P.O. Box 17155  
Austin, TX 78760-7155  
Phone: 800-531-5112  
Fax: (512) 447-7444

National Live Trap Corporation  
P.O. Box 302  
Tomahawk, WI 54487  
Phone: (715) 453-2249  
Fax: (715) 453-4705

National Netting, Inc.  
6325-C McDonough Drive  
Norcross, GA 30093  
Phone: 800-233-7896  
Fax: (404) 441-9260

Nature's All Natural Products, Inc.  
Houlton Road, P.O. Box 378  
Island Falls, ME 04747-0378  
Phone: (612) 548-3444  
Fax: (612) 548-3444

Nevada State Department of Agriculture  
Division of Agriculture  
350 Capitol Hill Avenue  
Reno, NV 89502  
Phone: (702) 688-1180  
Fax: (702) 688-1178

New Jersey Fireworks Company  
P.O. Box 118  
Vineland, NJ 08360  
Phone: (609) 692-8030

Nichols Net & Twine Co., Inc.  
2200 Highway 111  
Granite City, IL 62040  
Phone: (618) 797-0211  
Fax: (618) 797-0212

Nixalite of American, Inc.  
1025 16th Avenue, P.O. Box 727  
East Moline, IL 61244  
Phone: 800-624-1189  
Phone: (309) 755-8771  
Fax: (309) 755-0077

Norplex, Inc.  
111 3rd Street NW  
Auburn, WA 98002  
Phone: (206) 735-3431

Nortech Forest Products, Inc.  
7600 West 27th Street, Suite B-11  
St. Louis Park, MN 55426  
Phone: 800-323-3396  
Phone: (612) 922-2520  
Fax: (612) 922-3865

Nott Manufacturing Company, Inc.  
P.O. Box 685  
Pleasant Valley, NY 12569  
Phone: (914) 635-3243

Nylon Net Company  
615 East Bodley Avenue  
P.O. Box 592  
Memphis, TN 38101  
Phone: 800-238-7529  
Phone: (901) 774-1500  
Fax: (901) 775-5374

O'Gorman Enterprises  
P.O. Box 491  
Broadus, MT 59317  
Phone: (406) 436-2234

Orchard Equipment and Supply Company  
P.O. Box 540, Route 116  
Conway, MA 01341  
Phone: 800-634-5557

Orchard Supply Company  
P.O. Box 956  
Sacramento, CA 95812-0956  
Phone: (916) 446-7821

Pace International, Ltd.  
P.O. Box 558  
500 7th Avenue South  
Kirkland, WA 98083  
Phone: (206) 827-8711  
Fax: (206) 822-8261

Pace International, Ltd.  
Leffingwell Division  
111 South Berry Street, P.O. Box 1880  
Brea, CA 92621  
Phone: (714) 529-3973  
Fax: (714) 671-2138

Pacific Western Container  
1535 East Edinger Avenue  
Santa Ana, CA 92705  
Phone: (714) 547-9266  
Fax: (714) 953-9270

The Perry Company  
P.O. Box 7187  
Waco, TX 76714  
Phone: 800-792-3246  
Phone: (817) 756-2137  
Fax: (817) 756-2166

Pest Control Supplies  
1700 Liberty Street  
P.O. Box 025665  
Kansas City, MO 64102  
Phone: 800-821-5689  
Phone: (816) 421-4696  
Fax: (816) 472-0966

Pestcon Systems, Inc.  
5611 Capital Center Drive, Suite 302  
Raleigh, NC 27606  
Phone: 800-548-2778  
Phone: (919) 859-2500  
Fax: (919) 859-2155

Petersen Seeds, Inc.  
204 South Elm Street  
Gordon, NE 69343-1614  
Phone: (308) 282-1523

Pioneer Wildlife Traps  
2909 NE Alberta Street  
Portland, OR 97211  
Phone: (503) 249-2935

Pisces Industries  
P.O. Box 576407  
Modesto, CA 95355  
Phone: (209) 578-5502  
Fax: (209) 274-4723

Plant Pro-Tec, Inc.  
P.O. Box 902  
Palo Cedro, CA 96073  
Phone: (916) 547-5450  
Fax: (916) 547-5450

PMC Specialties Group, Inc.  
RJ Advantage, Inc.  
501 Murray Road  
Cincinnati, OH 45217  
Phone: (513) 482-7370  
Fax: (513) 482-7377

Pocatello Supply Depot  
USDA-APHIS, Animal Damage Control  
238 East Dillon  
Pocatello, ID 83201  
Phone: (208) 236-6920  
Fax: (208) 236-6922

Premier Fence Systems  
P.O. Box 89  
Washington, IA 52353  
Phone: 800-282-6631  
Fax: (319) 653-6304

Prosoco, Inc.  
P.O. Box 171677  
Kansas City, KS 66117  
Phone: 800-255-4255  
Phone: (913) 281-2700  
Fax: (913) 281-4385

P.S. Olt Company  
South 5th Street, Drawer 550  
Pekin, IL 61554  
Phone: (309) 348-3633  
Fax: (309) 348-3300

Purina Mills, Inc.  
P.O. Box 66812, 1401 South Hanley Road  
St. Louis, MO 63144  
Phone: (304) 768-4100  
Fax: (304) 768-4143

P-W Manufacturing Company  
610 High Street  
Henryetta, OK 74437  
Phone: (918) 652-4981



Qual-Line Fence Corporation  
801 South Division  
Waunakee, WI 53597  
Phone: (608) 849-4654

Quinn Manufacturing Company  
44201 Chapman Road  
Anza, CA 92539  
Phone: (909) 763-4590

Ranchers Supply, Inc.  
P.O. Box 725  
Alpine, TX 79831  
Phone: (915) 837-3630

RCO, Inc.  
P.O. Box 446  
Junction City, OR 97448  
Phone: 800-214-8160  
Fax: (503) 995-8160

Reed-Joseph International Company  
232 Main Street, P.O. Box 894  
Greenville, MS 38701  
Phone: 800-647-5554  
Fax: (601) 335-5822

Research Products Company  
Division of McShares, Inc.  
P.O. Box 1460  
Salina, KS 67402-1460  
Phone: (913) 825-2181  
Fax: (913) 825-8908

Rid-A-Bird, Inc.  
P.O. Box 436  
Wilton, IA 52778  
Phone: 800-432-4737  
Fax: (319) 732-3970

Rocky Mountain Wildlife Products  
4620 Moccasin Circle, P.O. Box 999  
Laporte, CO 80535  
Phone: (303) 484-2768

Rodent Control Systems, Inc.  
McKee Engineering Laboratories  
7207 Envoy Court, P.O. Box 561544  
Dallas, TX 75356  
Phone: (214) 638-5324  
Fax: (214) 638-5329

Roxide International, Inc.  
P.O. Box 249  
New Rochelle, NY 10802  
Phone: 800-431-5500  
Phone: (914) 235-5300  
Fax: (914) 235-5328

R-P Outdoors  
505 Polk Street, P.O. Box 1170  
Mansfield, LA 71052  
Phone: 800-762-2706  
Fax: (318) 872-8824

Rue R. Elston, Inc.  
706 North Weber  
Sioux Falls, SD 57102  
Phone: (605) 336-7716

Safe Shop Tools  
P.O. Box 4200  
Missoula, MT 59806  
Phone: 800-327-7639  
Fax: (406) 721-3545

Safeguard Products, Inc.  
P.O. Box 8  
New Holland, PA 17557  
Phone: 800-433-1819  
Phone: (717) 354-4586

Safety Live Trap  
P.O. Box 1226  
Townsend, MT 59644  
Phone: (406) 266-5590

Seabright Enterprises, Ltd.  
4067 Watts Street  
Emeryville, CA 94608-3604  
Phone: 800-284-7363  
Fax: (510) 654-7982

Sebesta Bait Mixing Plant  
P.O. Box 306  
Mitchell, SD 57301  
Phone: (605) 996-6131

Signal Broadcast Products, Inc.  
2314 Broadway  
Denver, CO 80205-2115  
Phone: (303) 296-1838  
Fax: (303) 296-1838

Sinco, Inc.  
P.O. Box 361  
East Hampton, CT 06424  
Phone: 800-243-6753  
Phone: (203) 267-2545

Smith & Hawken  
35 Corte Madera  
Mill Valley, CA 94941  
Phone: (415) 381-1800

The Snare Shop  
Rural Route 2, P.O. Box 147  
Carroll, IA 51401  
Phone: (712) 822-5318

Solvit, Inc.  
7001 Raywood Road  
Madison, WI 53713  
Phone: (608) 222-8624  
Fax: (608) 222-8733

South Dakota Department of Agriculture  
Rodent Control Fund  
Anderson Building  
Pierre, SD 57501  
Phone: (605) 773-3724

Southwest Chemical Company  
P.O. Box 1921  
Lompoc, CA 93438  
Phone: (805) 735-6156  
Fax: (805) 735-3629

Southwest Power Fence  
26321 Highway 281 North  
San Antonio, TX 78260  
Phone: 800-221-0178  
Phone: (210) 438-4600  
Fax: (210) 438-4604

Specialty Ag Equipment  
P.O. Box 1227, 344 East Dinuba Avenue  
Reedley, CA 93654  
Phone: 800-223-9799  
Phone: (209) 638-3631  
Fax: (209) 638-4710

Stendal Products, Inc.  
986 East Laurel Road  
Bellingham, WA 98226  
Phone: (360) 398-2353

Sterling Fur and Tool Company  
11268 Frick Road  
Sterling, OH 44276  
Phone: (216) 939-3763  
Fax: (216) 939-5135

Stoneco, Inc.  
P.O. Box 765  
Trinidad, CO 88082  
Phone: (719) 846-2853  
Fax: (719) 846-7700

Strieter Corporation  
2100 18th Avenue  
Rock Island, IL 61201-3611  
Phone: (309) 794-9800  
Fax: (309) 788-5646

Sudbury Laboratories, Inc.  
Division of Farnam Companies, Inc.  
P.O. Box 34820  
Phoenix, AZ 85067-4820  
Phone: 800-825-2555  
Phone: (602) 285-1660  
Fax: (602) 285-1803

Sullivan's Sure Catch Traps  
2324 South Patterson  
Valdosta, GA 31601  
Phone: (912) 242-1677

Sutton Ag Enterprises, Inc.  
746 Vertin Avenue  
Salinas, CA 93901  
Phone: (408) 422-9693  
Fax: (408) 422-4201

Techfence-Advanced Farm Systems  
Route 1, P.O. Box 364  
Bradford, ME 04410  
Phone: (207) 327-1237

Ted Dodge Service  
7604 Michel Road  
Mountain Ranch, CA 95246  
Phone: (209) 754-1216

Teiso Kasei Company  
350 South Figueroa Street, Suite 350  
Los Angeles, CA 90071  
Phone: (213) 608-4389

Tenax Corporation  
4800 East Monument Street  
Baltimore, MD 21205  
Phone: 800-356-8495  
Phone: (410) 522-7000  
Fax: (410) 522-7015

The Tensar Corporation  
1210 Citizens Parkway  
Morrow, GA 30260  
Phone: 800-292-4457  
Phone: (404) 968-3255  
Fax: (404) 961-8239

Tomahawk Live Trap Company  
P.O. Box 323  
Tomahawk, WI 54487  
Phone: 800-27-A-TRAP  
Phone: (715) 453-3550  
Fax: (715) 453-4326

Tomko Enterprises, Inc.  
180 Merritt Pond Road  
Riverhead, NY 11901  
Phone: (516) 727-3932

Topo Manufacturing & Welding Co.  
719 North C Street  
Imperial, CA 92251  
Phone: (619) 355-1556

Treessentials Company  
Riverview Station, P.O. Box 7097  
St. Paul, MN 55107  
Phone: 800-248-8239  
Fax: (612) 228-0554

Tripp-Lite Manufacturing Company  
500 North Orleans  
Chicago, IL 60610  
Phone: (312) 329-1777

Trucall  
9527 South Deer Creek Canyon Road  
Littleton, CO 80127  
Phone: (303) 697-4572  
Fax: (303) 697-4572

Twin Mountain Supply Company  
P.O. Box 2240  
San Angelo, TX 76902  
Phone: 800-527-0990  
Phone: (915) 944-8661  
Fax: (915) 949-2047

United Textile Corporation  
20 Universal Pl.  
Carlstat, NJ 07072  
Phone: (201) 993-9315  
Fax: (201) 993-1378

VanWaters and Rogers  
P.O. Box 34325  
Seattle, WA 98124-1325  
Phone: (206) 889-3400  
Fax: (206) 889-4100

Wald & Company  
208 Broadway  
Kansas City, MO 64105  
Phone: (806) 842-9299

Washington Forest Protection Association  
711 Capitol Way, Evergreen Plaza  
Building  
Olympia, WA 98501  
Phone: (206) 352-1500  
Fax: (206) 352-4621

Waterford Corporation  
404 North Link Lane, P.O. Box 1513  
Fort Collins, CO 80524  
Phone: 800-525-4952  
Phone: (303) 482-0911  
Fax: (303) 482-0934

Weitech, Inc.  
P.O. Box 1659, 310 Barclay Way  
Sisters, OR 97759  
Phone: 800-343-2659  
Phone: (503) 549-0205  
Fax: (503) 549-8154

West Virginia Fence Corporation  
U.S. Route 219  
Lindside, WV 24951  
Phone: 800-356-5458  
Phone: (304) 753-4387  
Fax: (304) 753-4827

Western Alfalfa  
P.O. Box 186  
Herndon, KS 67739  
Phone: (913) 322-5515

Western Fireworks Company  
2542 SE 13th Avenue  
Canby, OR 97013  
Phone: (503) 266-7770

Wickenkamp Live Trap Manufacturing  
Route 2  
Hedrick, IA 52563  
Phone: (515) 661-2156

Wilbur-Ellis Company  
191 West Shaw Avenue, Suite 107  
P.O. Box 1286  
Fresno, CA 98704-2876  
Phone: 800-937-3446  
Phone: (209) 226-1934  
Fax: (209) 226-7630

Wilco Distributors, Inc.  
1215 West Laurel Avenue  
P.O. Box 291  
Lompoc, CA 93436  
Phone: (805) 735-2476  
Fax: (805) 735-3629

Wildlife Control Technology, Inc.  
2501 North Sunnyside, #103  
Fresno, CA 93727  
Phone: 800-235-0262  
Phone: (209) 294-0262  
Fax: (209) 294-0632

Windear Manufacturing Company  
119 Parkway North, P.O. Box 415  
Lanesboro, MN 55949  
Phone: 800-726-5030  
Phone: (507) 467-3714

Woodstream Corporation  
69 North Locust  
Lititz, PA 17543-0327  
Phone: (717) 626-2125

York Distributors  
120 Express Street  
Plainview, NY 11803  
Phone: 800-645-6007  
Phone: (516) 932-0600  
Fax: (516) 932-4316

Z.A. Macabee Gopher Trap Company  
110 Loma Alta Avenue  
Los Gatos, CA 95032-6222  
Phone: (408) 354-4158

# Bibliography

- Andelt, W.F., D.L. Baker, and K.P. Burnham. 1991. Relative preference of captive cow elk for repellent-treated diets. *J. Wildl. Manage.* 56:164-173.
- Andelt, W.F., K.P. Burnham, and J.A. Manning. 1991. Relative effectiveness of repellents for reducing mule deer damage. *J. Wildl. Manage.* 55:341-347.
- Askam, L.R. 1992. Voles. pages 187-204 in H.C. Black (ed.), *Silvicultural approaches to animal damage management in Pacific Northwest Forests*. USDA Forest Service Technical Report PNW-GTR-287. Portland, Oregon.
- Baker, R.O., R.M. Timm, and G.R. Bodman. 1994. Rodent-proof construction and exclusion methods. pages B137-B150 in S.E. Hygnstrom, R.M. Timm, and G.E. Larson (eds.), *Prevention and Control of Wildlife Damage*. University of Nebraska Cooperative Extension. Lincoln, Nebraska.
- Bittner, S.L. and O.J. Rongstad. 1982. Snowshoe hare and allies. pages 146-163 in J.A. Chapman and G.A. Feldhamer (eds.), *Wild Mammals of North America*. Johns Hopkins University Press, Baltimore, Maryland.
- Black, H.C.. 1994. *Animal Damage Management Handbook*. USDA Forest Service, Pacific Northwest Res. Sta. PNW-GTR-332. 236 pp.
- Booth, T.W. 1994. Bird dispersal techniques. pages E19-E23 in S.E. Hygnstrom, R.M. Timm, and G.E. Larson (eds.), *Prevention and Control of Wildlife Damage*. University of Nebraska Cooperative Extension. Lincoln, Nebraska.
- Cafferata, S.L. 1992. Mountain beaver. pages 231-252 in H.C. Black (ed.), *Silvicultural approaches to animal damage management in Pacific Northwest Forests*. USDA Forest Service Technical Report PNW-GTR-287. Portland, Oregon.
- Campbell, D.L. 1994. Mountain beaver. pages B53-B60 in S.E. Hygnstrom, R.M. Timm, and G.E. Larson (eds.), *Prevention and Control of Wildlife Damage*. University of Nebraska Cooperative Extension. Lincoln, Nebraska.
- Campbell, D.L., and J. Evans. 1988. Recent approaches to controlling mountain beavers (*Apodonta rufa*) in Pacific Northwest Forests. *Proc. Vertebr. Pest Conf.* 13:183-187.
- Case, R.M., and B.A. Jasch. 1994. Pocket gophers. pages B117-B129 in S.E. Hygnstrom, R.M. Timm, and G.E. Larson (eds.), *Prevention and Control of Wildlife Damage*. University of Nebraska Cooperative Extension. Lincoln, Nebraska.
- Chase, J.D., W.E. Howard, and J.T. Roseberry. 1982. Pocket gophers. pages 239-255 in J.A. Chapman and G.A. Feldhamer (eds.), *Wild Mammals of North America*. Johns Hopkins University Press, Baltimore, Maryland.
- Clark, J. 1984. Vole control in field crops. *Proc. Vertebr. Pest Conf.* 11:5-6.
- Conover, M.R. 1984. Effectiveness of repellents in reducing deer damage in nurseries. *Wildl. Soc. Bull.* 12:399-404.
- Crouch, G.L. 1976. Deer and reforestation in the Pacific Northwest. *Proc. Vertebr. Pest Conf.* 7:298-301.
- Craven, S.R. 1994. Cottontail rabbits. pages D75-D80 in S.E. Hygnstrom, R.M. Timm, and G.E. Larson (eds.), *Prevention and Control of Wildlife Damage*. University of Nebraska Cooperative Extension. Lincoln, Nebraska.
- Craven, S.R., and S.E. Hygnstrom. 1994. Deer. pages D25-D40 in S.E. Hygnstrom, R.M. Timm, and G.E. Larson (eds.), *Prevention and Control of Wildlife Damage*. University of Nebraska Cooperative Extension. Lincoln, Nebraska.
- Crouch, G.L. 1986. Pocket gopher damage to conifers in western forests: a historical and current perspective on the problem and its control. *Proc. Vertebr. Pest Conf.* 12:196-198.
- deCalesta, D.S., and G.W. Witmer. 1994. Elk. pages D41-D50 in S.E. Hygnstrom, R.M. Timm, and G.E. Larson (eds.), *Prevention and Control of Wildlife Damage*. University of Nebraska Cooperative Extension. Lincoln, Nebraska.
- Dodge, W.E., and J.E. Borecco. 1992. Porcupines. pages 253-270 in H.C. Black (ed.), *Silvicultural Approaches to Animal Damage Management in Pacific Northwest Forests*. USDA Forest Service Technical Report PNW-GTR-287. Portland, Oregon.
- Dodge, W.E. 1982. Porcupine. pages 355-366 in J.A. Chapman and G.A. Feldhamer (eds.), *Wild Mammals of North America*. Johns Hopkins University Press, Baltimore, Maryland.
- Dolbeer, R.A., W.R. Holler, and D.W. Hawthorne. 1994. Identification and control of wildlife damage. pages 474-506 in T.A. Bookout (ed.), *Research and Management Techniques for Wildlife and Habitats*. The Wildlife Society, Bethesda, Maryland.
- Feldhamer, G.A., and J.A. Rochelle. 1982. Mountain beaver. pages 167-175 in J.A. Chapman and G.A. Feldhamer (eds.), *Wild Mammals of North America*. Johns Hopkins University Press, Baltimore, Maryland.
- Giusti, G.A., R.H. Schmidt, R.M. Timm, J.E. Borecco, and T.P. Sullivan. pages 289-308 in H.C. Black (ed.), *Silvicultural Approaches to Animal Damage Management in Pacific Northwest Forests*. USDA Forest Service Technical Report PNW-GTR-287. Portland, Oregon.
- Harris, M.T., W.L. Palmer, and J.L. George. 1983. Preliminary screening of white-tailed deer repellents. *J. Wildl. Manage.* 47:516-519.
- Hegdal, P.L., and T.A. Gatz. 1976. Hazards to wildlife associated with underground strychnine baiting for pocket gophers. *Proc. Vertebr. Pest Conf.* 7:158-266.
- Hesselton, W.T., and R.M. Hesselton. White-tailed deer. pages 878-901 in J.A. Chapman and G.A. Feldhamer (eds.), *Wild Mammals of North America*. Johns Hopkins University Press, Baltimore, Maryland.
- Hill, E.P. 1982. Beaver. pages 256-281. in J.A. Chapman and G.A. Feldhamer (eds.), *Wild Mammals of North America*. Johns Hopkins University Press, Baltimore, Maryland.

- Hygnstrom, S.E. 1994. Black bears. pages C5-C15 in S.E. Hygnstrom, R.M. Timm, and G.E. Larson (eds.), *Prevention and Control of Wildlife Damage*. University of Nebraska Cooperative Extension. Lincoln, Nebraska.
- Hygnstrom, S.E., and S.R. Craven. 1988. Electric fences and commercial repellents for reducing deer damage in cornfields. *Wild. Soc. Bull.* 16:291-296.
- Hygnstrom, S.E., and D.J. Hafer. 1994. pages H1-H31 in S.E. Hygnstrom, R.M. Timm, and G.E. Larson (eds.), *Prevention and Control of Wildlife Damage*. University of Nebraska Cooperative Extension. Lincoln, Nebraska.
- Jackson, J.J. 1994. Tree squirrels. pages B171-B175 in S.E. Hygnstrom, R.M. Timm, and G.E. Larson (eds.), *Prevention and Control of Wildlife Damage*. University of Nebraska Cooperative Extension. Lincoln, Nebraska.
- Jacobs, W.W. 1994. Registered vertebrate pesticides. pages G1-G22 in S.E. Hygnstrom, R.M. Timm, and G.E. Larson (eds.), *Prevention and Control of Wildlife Damage*. University of Nebraska Cooperative Extension. Lincoln, Nebraska.
- Johnson, M.L., and S. Johnson. 1982. Voles. pages 326-354 in J.A. Chapman and G.A. Feldhamer (eds.), *Wild Mammals of North America*. Johns Hopkins University Press, Baltimore, Maryland.
- Karsky, R. 1988. *Fences*. USDA Forest Service—Missoula Technology and Development Center, Missoula, Montana. 1988-594-194/80139. 210 pp.
- Knapp, W.H., and J.D. Brodie. 1992. The process of managing animal damage. pages 365-376 in H.C. Black (ed.), *Silvicultural Approaches to Animal Damage Management in Pacific Northwest Forests*. USDA Forest Service Technical Report PNW-GTR-287. Portland, Oregon.
- Lund, M. 1988. Nonanticoagulant rodenticides. pages 331-340 in I. Prakash (ed.), *Rodent Pest Management*. CRC Press Inc., Boca Raton, Florida.
- Lund, M. 1988. Anticoagulant rodenticides. pages 341-351 in I. Prakash (ed.), *Rodent Pest Management*. CRC Press Inc., Boca Raton, Florida.
- Mackie, R.J., K.L. Hamlin, and D.F. Pac. 1982. Mule Deer. pages 862-877 in J.A. Chapman and G.A. Feldhamer (eds.), *Wild Mammals of North America*. Johns Hopkins University Press, Baltimore, Maryland.
- Marsh, R.E. 1992. Reflections on current (1992) pocket gopher control in California. *Proc. Vertebr. Pest Conf.* 15:289-295.
- Marsh, R.E., and W.E. Howard. 1990. Vertebrate pests. pages 791-861 in A. Mallis (ed.), *Handbook of Pest Control*, 7th edition. Franzak and Foster Co., Cleveland, Ohio.
- Marsh, R.E., and R.W. Steele. 1992. Pocket gophers. pages 205-230 in H.C. Black (ed.), *Silvicultural Approaches to Animal Damage Management in Pacific Northwest Forests*. USDA Forest Service Technical Report PNW-GTR-287. Portland, Oregon.
- Miller, J.E., and G.K. Yarrow. 1994. Beavers. pages B1-B11 in S.E. Hygnstrom, R.M. Timm, and G.E. Larson (eds.), *Prevention and Control of Wildlife Damage*. University of Nebraska Cooperative Extension. Lincoln, Nebraska.
- O'Brien, J.M. 1994. Voles. pages B177-B182 in S.E. Hygnstrom, R.M. Timm, and G.E. Larson (eds.), *Prevention and Control of Wildlife Damage*. University of Nebraska Cooperative Extension. Lincoln, Nebraska.
- Palmer, W.L., R.G. Wingard, and J.L. George. 1983. Evaluation of white-tailed deer repellents. *Wildl. Soc. Bull.* 11:164-166.
- Pauls, D.T. 1986. Protection with Vexar cylinders from damage by meadow voles of tree and shrub seedlings in northeastern Alberta. *Proc. Vertebr. Pest Conf.* 12:199-204.
- Peek, J.M. 1982. Elk. pages 851-861 in J.A. Chapman and G.A. Feldhamer (eds.), *Wild Mammals of North America*. Johns Hopkins University Press, Baltimore, Maryland.
- Pelton, M.R. 1982. Black bear. pages 504-514 in J.A. Chapman, and G.A. Feldhamer (eds.), *Wild Mammals of North America*. Johns Hopkins University Press, Baltimore, Maryland.
- Rochelle, J.A. 1992. Deer and elk. pages 333-350 in H.C. Black (ed.), *Silvicultural Approaches to Animal Damage Management in Pacific Northwest Forests*. USDA Forest Service Technical Report PNW-GTR-287. Portland, Oregon.
- Schmidt, W.C., and M. Gourley. 1992. pages 309-332 in H.C. Black (ed.), *Silvicultural Approaches to Animal Damage Management in Pacific Northwest Forests*. USDA Forest Service Technical Report PNW-GTR-287. Portland, Oregon.
- Schmitz, S.D. 1994. Porcupines. pages B81-B83 in S.E. Hygnstrom, R.M. Timm, and G.E. Larson (eds.), *Prevention and Control of Wildlife Damage*. University of Nebraska Cooperative Extension. Lincoln, Nebraska.
- Sullivan, T.P. 1992. Tree squirrels, woodrats, and beaver. pages 271-288 in H.C. Black (ed.), *Silvicultural Approaches to Animal Damage Management in Pacific Northwest Forests*. USDA Forest Service Technical Report PNW-GTR-287. Portland, Oregon.
- Swihart, R.K., and M.R. Conover. 1990. Reducing deer damage to yews and apple trees: testing Big Game Repellents, Ro-pel, and soap as repellents. *Wildl. Soc. Bull.* 18:156-162.
- Weaver, K.M., D.H. Arner, C. Mason, and J.J. Hartely. 1985. A guide to using snares for beaver captures. *South. J. Appl. For.* 9:141-146.

**For further technical information,  
please contact:**

Dr. Dale Nolte, Wildlife Biologist  
USDA-APHIS, Animal Damage Control  
Denver Wildlife Research Center  
9701 Blomberg Street  
Olympia, WA 98512  
Phone: (360) 664-3341  
Fax: (360) 664-3289

**For additional copies of this  
publication, contact:**

USDA Forest Service, MTDC  
Building 1, Fort Missoula  
Missoula, MT 59801  
Phone: (406) 329-3900  
Fax: (406) 329-3719

**Library Card**

Nolte, Dale L.; Otto, Ivy J. 1996.  
Materials and supplies for management  
of wildlife damage to trees. Tech. Rep.  
9624-2808-MTDC. Missoula, MT: USDA  
Forest Service, Missoula Technology  
and Development Center. 48 p.

Summarizes ways to control damage to  
trees by wildlife. Includes tips for  
identifying problems caused by  
mountain beavers, beavers, elk,  
porcupines, snowshoe hares, voles,  
deer, pocket gophers, and black bears.  
Provides information on physical  
deterrents, traps, toxicants, repellents,  
frightening devices, and alternative  
forages. Lists more than 200  
companies supplying materials.

Keywords: repellents, traps, toxicants,  
reforestation.